

## PROBLEM 10.1

Two identical spheres, each of mass  $m$ , are suspended from a horizontal ceiling by two strings of length  $l$ . The strings are attached to the ceiling at a distance  $2l$  apart. The spheres are released from rest at an angle  $\theta_0$  to the vertical. Determine the velocity of each sphere at the instant the strings are vertical.

Hint: Use the principle of conservation of energy.

Assume that the strings are inextensible and that the spheres do not collide.

Assume that the spheres are released from rest at an angle  $\theta_0$  to the vertical.

Assume that the strings are inextensible and that the spheres do not collide.

Assume that the spheres are released from rest at an angle  $\theta_0$  to the vertical.

Assume that the strings are inextensible and that the spheres do not collide.

Assume that the spheres are released from rest at an angle  $\theta_0$  to the vertical.

Assume that the strings are inextensible and that the spheres do not collide.

Assume that the spheres are released from rest at an angle  $\theta_0$  to the vertical.

Assume that the strings are inextensible and that the spheres do not collide.

Assume that the spheres are released from rest at an angle  $\theta_0$  to the vertical.

Assume that the strings are inextensible and that the spheres do not collide.

Assume that the spheres are released from rest at an angle  $\theta_0$  to the vertical.

Assume that the strings are inextensible and that the spheres do not collide.

Assume that the spheres are released from rest at an angle  $\theta_0$  to the vertical.

