

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On admission, the patient is found to have a serum sodium of 125 mEq/L, serum glucose of 100 mg/dL, and serum urea nitrogen of 18 mg/dL. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

Test	Result
Serum sodium	125 mEq/L
Serum glucose	100 mg/dL
Serum urea nitrogen	18 mg/dL
Serum creatinine	1.2 mg/dL
Serum albumin	3.5 g/dL
Serum total bilirubin	2.5 mg/dL
Serum aspartate aminotransferase (AST)	150 U/L
Serum alanine aminotransferase (ALT)	100 U/L
Serum gamma-glutamyl transaminase (GGT)	250 U/L
Serum lactate dehydrogenase (LDH)	1000 U/L
Serum ferritin	100 ng/mL
Serum ferritin (reference range)	50-150 ng/mL

2. A 65-year-old male patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On admission, the patient is found to have a serum sodium of 125 mEq/L, serum glucose of 100 mg/dL, and serum urea nitrogen of 18 mg/dL. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

Test	Result
Serum sodium	125 mEq/L
Serum glucose	100 mg/dL
Serum urea nitrogen	18 mg/dL
Serum creatinine	1.2 mg/dL
Serum albumin	3.5 g/dL
Serum total bilirubin	2.5 mg/dL
Serum aspartate aminotransferase (AST)	150 U/L
Serum alanine aminotransferase (ALT)	100 U/L
Serum gamma-glutamyl transaminase (GGT)	250 U/L
Serum lactate dehydrogenase (LDH)	1000 U/L
Serum ferritin	100 ng/mL
Serum ferritin (reference range)	50-150 ng/mL

3. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On admission, the patient is found to have a serum sodium of 125 mEq/L, serum glucose of 100 mg/dL, and serum urea nitrogen of 18 mg/dL. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

ANSWER



4. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On admission, the patient is found to have a serum sodium of 125 mEq/L, serum glucose of 100 mg/dL, and serum urea nitrogen of 18 mg/dL. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

Test	Result
Serum sodium	125 mEq/L
Serum glucose	100 mg/dL
Serum urea nitrogen	18 mg/dL
Serum creatinine	1.2 mg/dL
Serum albumin	3.5 g/dL
Serum total bilirubin	2.5 mg/dL
Serum aspartate aminotransferase (AST)	150 U/L
Serum alanine aminotransferase (ALT)	100 U/L
Serum gamma-glutamyl transaminase (GGT)	250 U/L
Serum lactate dehydrogenase (LDH)	1000 U/L
Serum ferritin	100 ng/mL
Serum ferritin (reference range)	50-150 ng/mL

5. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On admission, the patient is found to have a serum sodium of 125 mEq/L, serum glucose of 100 mg/dL, and serum urea nitrogen of 18 mg/dL. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows: