

QUESTION

1. A patient with a long history of chronic kidney disease (CKD) is being treated with a loop diuretic. The patient's serum potassium level is 2.8 mEq/L. The patient is also taking a beta-blocker and a calcium channel blocker. The patient's blood pressure is 150/90 mmHg. The patient is also taking a proton pump inhibitor (PPI) and a statin. The patient's creatinine level is 2.5 mg/dL. The patient is also taking a diuretic. The patient is also taking a beta-blocker and a calcium channel blocker. The patient's blood pressure is 150/90 mmHg. The patient is also taking a proton pump inhibitor (PPI) and a statin. The patient's creatinine level is 2.5 mg/dL. The patient is also taking a diuretic.

Parameter	Value
Serum Potassium	2.8 mEq/L
Blood Pressure	150/90 mmHg
Creatinine	2.5 mg/dL

2. A patient with a long history of chronic kidney disease (CKD) is being treated with a loop diuretic. The patient's serum potassium level is 2.8 mEq/L. The patient is also taking a beta-blocker and a calcium channel blocker. The patient's blood pressure is 150/90 mmHg. The patient is also taking a proton pump inhibitor (PPI) and a statin. The patient's creatinine level is 2.5 mg/dL. The patient is also taking a diuretic. The patient is also taking a beta-blocker and a calcium channel blocker. The patient's blood pressure is 150/90 mmHg. The patient is also taking a proton pump inhibitor (PPI) and a statin. The patient's creatinine level is 2.5 mg/dL. The patient is also taking a diuretic.

ANSWER



3. A patient with a long history of chronic kidney disease (CKD) is being treated with a loop diuretic. The patient's serum potassium level is 2.8 mEq/L. The patient is also taking a beta-blocker and a calcium channel blocker. The patient's blood pressure is 150/90 mmHg. The patient is also taking a proton pump inhibitor (PPI) and a statin. The patient's creatinine level is 2.5 mg/dL. The patient is also taking a diuretic. The patient is also taking a beta-blocker and a calcium channel blocker. The patient's blood pressure is 150/90 mmHg. The patient is also taking a proton pump inhibitor (PPI) and a statin. The patient's creatinine level is 2.5 mg/dL. The patient is also taking a diuretic.