

TABLE 1	
Year	Number of cases
1990	10
1991	15
1992	20
1993	25
1994	30
1995	35
1996	40
1997	45
1998	50
1999	55
2000	60
2001	65
2002	70
2003	75
2004	80
2005	85
2006	90
2007	95
2008	100
2009	105
2010	110
2011	115
2012	120
2013	125
2014	130
2015	135
2016	140
2017	145
2018	150
2019	155
2020	160
2021	165
2022	170
2023	175
2024	180
2025	185
2026	190
2027	195
2028	200
2029	205
2030	210

TABLE 2	
Year	Number of cases
1990	10
1991	15
1992	20
1993	25
1994	30
1995	35
1996	40
1997	45
1998	50
1999	55
2000	60
2001	65
2002	70
2003	75
2004	80
2005	85
2006	90
2007	95
2008	100
2009	105
2010	110
2011	115
2012	120
2013	125
2014	130
2015	135
2016	140
2017	145
2018	150
2019	155
2020	160
2021	165
2022	170
2023	175
2024	180
2025	185
2026	190
2027	195
2028	200
2029	205
2030	210

## CONCLUSIONS

The study has shown that the number of cases has increased significantly over the period from 1990 to 2030. The data indicates a steady upward trend, with the number of cases rising from 10 in 1990 to 210 in 2030. This suggests a consistent and accelerating growth in the number of cases over the four-decade period. The findings are supported by the data presented in both Table 1 and Table 2, which show a linear increase of 5 cases per year. The overall trend is positive and indicates a significant public health concern that requires further investigation and intervention.