



Specifications

Color	Yellow
Appearance	Clear
Fragrance	Lemon
Formula Type	Liquid
Dilution Ratio	1:128
PH	12.00 - 13.00
Non-Volatile Matter	19.50 - 21.50%
EPA Registration #	6836-77-1658

Safety

See safety datasheet (if applicable) and product label for safety information, handling and proper use.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected. **Preparation of Disinfectant Use-Solution:** Add 1 oz. per gallon of water (1:128). For heavy duty use, add 4 oz. per gallon of water (1:32). Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Use 16 oz. per gallon of water to kill Feline Panleukopenia Virus. **DISINFECTION / VIRUCIDAL* / FUNGICIDAL / MOLD AND MILDEW CONTROL DIRECTIONS:** Apply use-solution to hard, non-porous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge, sprayer or by immersion. Treated surfaces must remain visibly wet for 10 minutes. **For Influenza Virus Type A, Human Coronavirus and SARS-Cov-2,** treated surfaces must remain visibly wet for 1 minute. Wipe dry with a cloth, sponge or mop or allow to air dry. For visibly soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface. Do not breathe spray. Rinse all surfaces that come in contact with food

Q.T. Plus has been formulated to aid in the reduction of cross-contamination between treated hard, non-porous surfaces not only in hospitals, but in schools, institutions, and industry.

Effective in hard water up to 200 ppm hardness (calculated as CaCO₃) in the presence of 5% serum contamination. Q.T. Plus is a one-step Hospital use disinfectant, Bactericidal tested according to the current AOAC Use-Dilution Test Method and Virucidal* tested according to the virucidal qualification modified in the presence of 200 ppm hard water plus 5% organic serum. 1 MINUTE CONTACT TIME: *SARS-Related Coronavirus 2 (SARS-CoV-2) (cause of COVID-19).

Features & Benefits

Higher pH removes tough soils ordinary neutral pH disinfectants can not.

Mild lemon fragrance.

EPA registered disinfectant.

Effective against SARS-CoV-2, the virus that causes COVID-19 with a 3- minute contact time.

Item Number & Unit of Measure

Item Number	HIL0016909
Unit of Measure	55 gal Drum

such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes. **For fungicidal activity:** Q.T. Plus is an effective fungicide against *Trichophyton interdigitale* (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. Dilute at rate of 1 oz. per gallon in 200 ppm hard water and allow surface to remain visibly wet for 10 minutes. **For mold and mildew:** At 1 oz. per gallon, Q.T. Plus will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non-porous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

BACTERICIDAL STABILITY OF USE-

DILUTION: Tests confirm that the use solution of this product, when stored in a sealed container such as a spray bottle, remains effective for up to one month. If product becomes visibly dirty or contaminated, the use-dilution must be discarded and fresh product prepared. Always use clean, properly labeled containers when diluting this product. Bactericidal stability of the use-dilution does not apply to open containers such as buckets or pails. This product is not for use on medical device surfaces. ***KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

Special Instructions for Cleaning and Decontamination against HIV-1 or HBV or HCV on Surfaces / Objects Soiled With Blood / Body Fluids.

Personal Protection: Cleanup must always be done wearing protective gloves, gowns, masks

and eye protection. **Cleaning Procedure:** Blood and other body fluids containing HIV-1 or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of Q.T. Plus. **Contact Time:** Leave surface visibly wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV with a 1 oz. per gallon use-solution. Use a 10-minute contact time for disinfection against all other viruses, bacteria and fungi claimed. **Disposal of Infectious Material:** Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal. **For use as a One-Step Cleaner and Disinfectant by Electrostatic**

Spraying1:

Electrostatic spray application is for efficacy claims with a 10 minute contact time and 1:128 dilution. This application method is not to be used to treat surfaces for sanitization or fungicidal claims.

- Ensure bystanders and pets are not present in the room during application.
- Clean visibly dirty surfaces prior to spraying.
- Wear a N95 filtering facepiece respirator or half face respirator with a N95 filter and eye protection (goggles, face shield or safety glasses) when spraying.
- Select a spray nozzle that produces spray droplet particles with a volume median diameter (VMD) $\geq 80 \mu\text{m}$.
- Place the electrostatic spray function in the ON position for electrostatic spray models that have the functionality to toggle ON/OFF.
- Hold the sprayer nozzle 12 inches away from the target surface. Starting at the highest point and working down to the lowest point, spray target surfaces using a smooth methodical "S" pattern. To avoid leaving gaps in sprayed surfaces, overlap previously treated surfaces when moving onto subsequent sections.
- The target surface must remain visibly wet for 10 minutes. Re-apply product as necessary to keep surface visibly wet for the duration of the contact time. Wipe surfaces dry (or rinse) (or allow to air dry).

Q.T.® Plus Disinfectant Cleaner (continued)



· Allow the treated room to remain unoccupied for 15 minutes before reentering. **NON-ACID TOILET BOWL AND URINAL DISINFECTANT / CLEANER DIRECTIONS:** Remove visible filth prior to disinfection.

From Concentrate: Add 1 oz. to the bowl. Use 4 oz. to kill Adenovirus Type 5 and Type 7. Thoroughly brush or swab over exposed surfaces and under the rim with a cloth, mop or sponge. Allow to stand for 10 minutes and flush.

From Use Solution: Empty toilet bowl or urinal and apply a 1 oz. per gallon use-solution to exposed surfaces and under the rim with a cloth, mop, sponge or sprayer. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Brush or swab thoroughly, allow to stand for 10 minutes and flush.

For Heavy Duty Use: Empty toilet bowl or urinal and apply a 4 oz. per gallon use-solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly, allow to stand for 10 minutes and flush. **VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY/ZOOS/PET SHOPS /**

KENNELS DISINFECTION DIRECTIONS: For cleaning and disinfecting hard, non-porous surfaces: Equipment used for feeding or watering animals, utensils, instruments, cages, kennels, stables, catteries, pens, stalls and etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, pens, stalls and etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 1 oz. of **Q.T. Plus** per gallon of water for a period of 10 minutes. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Use 16 oz. per gallon of water to kill Feline Panleukopenia Virus. Wipe or allow to air dry. Immerse all animal halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure.

Thoroughly scrub all treated surfaces, then rinse all surfaces that come in contact with food, including equipment used for feeding or watering, with potable water before reuse. Ventilate buildings, animal enclosures, vehicles and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. **FARM PREMISE USE:** Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with a use-solution of 1 oz. of **Q.T. Plus** per gallon of water for a period of 10 minutes. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure. Ventilate buildings, cars, trucks, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse. **TO SANITIZE NON-FOOD CONTACT SURFACES:** Add 1 oz. of **Q.T. Plus** per 4.25 gallons of water to sanitize hard, non-porous surfaces. Treated surfaces must remain visibly wet for 60 seconds. Wipe dry with a sponge, mop or cloth or allow to air dry. Do not use in food preparation, food handling and food storage areas. This product is not to be used on glassware, dishware, cookware or utensils. **TO CLEAN AND DEODORIZE:** Mix 1 oz. of **Q.T. Plus** per gallon of water to clean and deodorize hard, non-porous, non-food surfaces. Apply using a cloth, mop, sponge or sprayer. Wipe or allow to air dry. Rinse all surfaces that come in contact with food such as countertops, exteriors of appliances, tables

Q.T.® Plus Disinfectant Cleaner (continued)



and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes.

Efficacy

BACTERICIDAL ACTIVITY: 10-minute contact time Pseudomonasaeruginosa, Staphylococcus aureus, Salmonella enterica, Acinetobacterbaumanni, Corynebacterium ammoniagenes, Campylobacter jejuni, Staphylococcus aureus - Community Associated Methicillin-Resistant - CA-MRSA (NRS384)(USA300), Staphylococcus aureus - Community Associated Methicillin-Resistant -CA-MRSA (NRS123) (USA400), Enterobacteraerogenes, Enterobacter cloacae NDM-1 positive, Enterococcus faecalis, Enterococcus faecalis-Vancomycin resistant (VRE), Escherichia coli, Escherichiacoli, New Delhi Metallo-Beta Lactamase (NDM-1), ESBL Escherichia coli, Klebsiella pneumoniae, Klebsiellapneumoniae - NDM-1 positive, Listeria monocytogenes, Legionellapneumophila, Burkholderia cepacia, Salmonella schottmuelleri, Salmonella typhi, Serratia marcescens, Shigelladysenteriae, Staphylococcus aureus Methicillin-resistant (MRSA), Staphylococcus aureus Multi-Drug Resistant (resistant to tetracycline (Tc), penicillin (Pc), streptomycin (Sm) and erythromycin (Em) and susceptible to chloramphenicol (Cm) *in vitro*), Staphylococcus aureus-Vancomycin Intermediate Resistant-(VISA), Streptococcus pyogenes, Vibrio cholerae. **VIRUCIDAL ACTIVITY: 10-minute** contact time *Hepatitis B Virus (HBV), *Hepatitis C Virus (HCV), *Herpes Simplex Virus Type 1, *Herpes Simplex Virus Type 2, *Norwalk Virus (Norovirus), *Rotavirus, *Respiratory Syncytial Virus (RSV), *SARS Associated Coronavirus, *Vaccinia. Fungi: Aspergillus niger, Trichophyton mentagrophytes (the Athlete's Foot Fungus) (A cause of Ringworm of the foot). Avian Infectious Bronchitis Virus, Avian Influenza Virus (H3N2), Avian Influenza Virus (H5N1), Canine Distemper Virus, Feline Calicivirus, Feline Panleukopenia Virus (16 oz. per gallon of water), Newcastle's Disease Virus, Pseudorabies Virus. **1-minute** contact time *SARS-Related Coronavirus 2 (cause of COVID-19), *HIV-1 (AIDS virus), *Human Coronavirus, *Influenza Virus Type A / Brazil, Influenza

Virus Type A/Hong Kong, *2013 Influenza A Virus (H7N9). **FUNGICIDAL ACTIVITY: 10-minute** contact time, Trichophyton interdigitale (the athlete's foot fungus).

HMIS Ratings

Flammability RTU	0
Health RTU	1
Reactivity RTU	0