



Hillyard Quick and Clean Q.T. 4 Disinfectant Wipes contain a hospital-grade quat-based formula. They are bleach-free, alcohol-free, and non-staining. They are ideal for use as a cleaner and disinfectant in Exercise Facilities, Health Clubs, Locker Rooms, Hospitals, Dental Offices, Nursing Homes, Day Care Centers, Institutional Kitchens, USDA-inspected food Processing Facilities, and many other environments. Disinfect hard, non-porous surfaces in 4 minutes and sanitize in 15 seconds. They also effectively kill mold and mildew and have a pleasant lemon fragrance.

Features & Benefits

Hospital-grade quat-based formula.

Bleach-free, alcohol-free, and non-staining, ensuring safety and suitability for diverse environments.

Disinfect hard surfaces in 4 minutes and sanitize in 15 seconds.

Kills mold and mildew.

Item Number & Unit of Measure

| | |
|-----------------|---------------------|
| Item Number | HIL01175 |
| Unit of Measure | 180 Wipes/Container |

Specifications

| | |
|--------------|-------|
| Color | White |
| Fragrance | Lemon |
| Formula Type | Wipes |

| | |
|--------------------|----------------|
| PH | > 7.0 - < 10.5 |
| EPA Registration # | 6836-340-1658 |

Safety

See safety datasheet (if applicable) and product label for safety information, handling and proper use.

Directions for Use

Do not use as a diaper wipe or for personal cleansing. This is not a baby wipe! Do not flush in toilet.

For plastic and painted surfaces, spot test on an inconspicuous area before use. This product will not harm most surfaces: including acrylic, sealed fiberglass and vinyl.

TO CLEAN AND DISINFECT/ VIRUCIDAL* DIRECTIONS: Wipe hard, nonporous surface with wipe until surface is visibly wet. Allow surface to remain visibly wet for 4 minutes. For Norwalk Virus, Feline Calicivirus, Norovirus and Rotavirus, treated surfaces must remain visibly wet for 10 minutes. For SARS-CoV-2, treated surfaces must remain visibly wet for 15 seconds. Let air dry. If surfaces are visibly dirty, clean first with another wipe before disinfecting. Discard used wipe in trash. Do not flush.

For mold and mildew stain control, this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Wipe surface and let it remain visibly wet for 4 minutes. Repeat treatment every seven days, or more often if new growth appears.

KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

Special Instructions for Cleaning and Decontamination against HIV-1 or HBV or HCV on Surfaces / Objects Soiled With Blood / Body Fluids. Personal protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection. **Cleaning procedure:** Blood and other body fluids containing HIV or HBV or HCV must be visibly cleaned from surfaces and objects before application of this product can be used for this purpose. **Contact Time:** Leave surface visibly wet for 1 minute for HIV-1. Use appropriate contact time as listed on label for all other

Q.T. 4 Disinfectant Wipes (continued)



viruses and bacteria claimed. **Disposal of infectious material:** Blood, body fluids, cleaning materials and clothing must be disposed of according to local regulations for infectious waste disposal.

| | |
|------------------|---|
| Flammability RTU | 0 |
| Health RTU | 0 |
| Reactivity RTU | 0 |

This product is not for use on medical device surfaces.
This product is for use on hard nonporous surfaces not conducive to treatment by immersion or excess liquid against Avian Influenza A virus.

SANITIZING, DEODORIZING: To sanitize, deodorize hard nonporous, non-food contact surfaces; wipe surface, using enough wipes to allow treated surface to remain visibly wet for 15 seconds for *Staphylococcus aureus* (ATCC 6538), *Klebsiella pneumoniae* (ATCC 4352), and *Campylobacter jejuni* (ATCC 29428). Let surface dry. For visibly soiled surfaces, clean visible soil first.

Efficacy

Effective against bacteria including *Pseudomonas aeruginosa* (*Pseudomonas*), *Salmonella enterica* (*Salmonella*), *Staphylococcus aureus* (*Staph*), *Acinetobacter baumannii*, *Brevibacterium ammoniagenes* (*Brevibacterium*), *Campylobacter jejuni* (*Campylobacter*), Community Associated Methicillin - Resistant *Staphylococcus aureus* - (CA-MRSA), *Enterobacter aerogenes*, *Enterococcus faecalis* (*Enterococcus*), *Enterococcus faecalis* - Vancomycin Resistant (VRE), *Escherichia coli* (*E. coli*), *Escherichia coli* O157:H7, ESBL *Escherichia coli* - (Extended spectrum beta-lactamase producing *E. coli*), *Klebsiella pneumoniae* (*Klebsiella*), *Legionella pneumophila*, *Listeria monocytogenes*, *Pseudomonas cepacia* (*Pseudomonas*), *Salmonella schottmuelleri* (*Salmonella*), *Salmonella typhi* (*Salmonella*), *Serratia marcescens* (*Serratia*), *Shigella dysenteriae* (*Shigella*), *Staphylococcus aureus* - Methicillin-Resistant (MRSA), *Staphylococcus aureus* - Multi-Drug Resistant (resistant to tetracycline [Tc], penicillin [Pc], streptomycin [Sm] and erythromycin [Em] and susceptible to chloramphenicol [Cm] in vitro), *Staphylococcus aureus*-Vancomycin Intermediate Resistant - (VISA), *Streptococcus pyogenes* (*Strep*), *Vibrio cholera*.

Effective against virus including: Herpes Simplex Virus Type 1, Herpes Simplex Virus Type 2, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), HIV-1 (AIDS Virus), Human Coronavirus, Influenza Type A Virus / Brazil Influenza, Influenza A Virus H1N1, Norwalk Virus (Norovirus)*, Respiratory Syncytial Virus (RSV), Rotavirus, SARS Associated Coronavirus (SARS) (cause of Severe Acute Respiratory Syndrome), and Vaccinia Virus (Pox Virus). *Treated surfaces must remain wet for 10 minutes.