

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, anorexia, and weakness. Physical examination reveals a 10% weight loss, a dry mucous membrane, and a heart rate of 100 beats per minute. Laboratory studies show a hemoglobin of 10 g/dL, a hematocrit of 30%, and a mean corpuscular volume of 80 fL. The patient's serum iron is 50 µg/dL, and the total iron-binding capacity is 300 µg/dL. The patient's serum ferritin is 100 µg/L. The patient's serum transferrin saturation is 16.7%. The patient's serum ferritin is 100 µg/L. The patient's serum transferrin saturation is 16.7%. The patient's serum ferritin is 100 µg/L. The patient's serum transferrin saturation is 16.7%.

ANSWER

1. The patient's laboratory studies are consistent with iron deficiency anemia. The patient's hemoglobin is 10 g/dL, and the hematocrit is 30%. The patient's mean corpuscular volume is 80 fL, which is consistent with a microcytic anemia. The patient's serum iron is 50 µg/dL, and the total iron-binding capacity is 300 µg/dL. The patient's serum ferritin is 100 µg/L. The patient's serum transferrin saturation is 16.7%.

2. The patient's laboratory studies are consistent with iron deficiency anemia. The patient's hemoglobin is 10 g/dL, and the hematocrit is 30%. The patient's mean corpuscular volume is 80 fL, which is consistent with a microcytic anemia. The patient's serum iron is 50 µg/dL, and the total iron-binding capacity is 300 µg/dL. The patient's serum ferritin is 100 µg/L. The patient's serum transferrin saturation is 16.7%.

QUESTION



3. The patient's laboratory studies are consistent with iron deficiency anemia. The patient's hemoglobin is 10 g/dL, and the hematocrit is 30%. The patient's mean corpuscular volume is 80 fL, which is consistent with a microcytic anemia. The patient's serum iron is 50 µg/dL, and the total iron-binding capacity is 300 µg/dL. The patient's serum ferritin is 100 µg/L. The patient's serum transferrin saturation is 16.7%.