



Formula Type	Foam
PH	11.82 - 12.82
EPA Registration #	706-65-1658

Safety

See safety datasheet (if applicable) and product label for safety information, handling and proper use.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. To Clean and Deodorize: Remove all gross filth or heavy soil from surfaces to be treated prior to using this product. Shake Well Before Each Use. Point valve opening toward area to be cleaned, press down firmly and spray evenly. Wipe clean with a dry cloth, turning frequently. This product effectively controls the growth of mold and mildew and the odors caused by them when applied to hard, non-porous surfaces. Apply by holding can upright 6-8 inches from surface to be treated. Spray until surface is completely wet. Let air dry. Repeat application at weekly intervals or when mildew growth appears. This product demonstrates effective fungicidal activity against pathogenic fungi, Trichophyton mentagrophytes, in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities and on clean hard, non-porous surfaces commonly contacted by bare feet. Follow disinfection directions. To Disinfect: After surface is cleaned, hold can upright 6-8 inches from surface to be treated. Spray until surface is thoroughly wet. Allow to dry without wiping or allow spray to contact treated surface for a minimum of ten minutes prior to wiping. When used as a disinfectant on food contact surfaces such as counter tops, tables and appliances, thoroughly rinse surfaces with potable water. Do not use on utensils, dishes or glasses. Kills HIV on Pre-Cleaned Environmental Surfaces Previously Soiled with Blood/Body Fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV) (associated with AIDS): Special Instructions for Cleaning and Decontamination Against HIV on Surfaces/Objects Soiled with Blood/Body Fluids: Personal Protection: When handling items soiled with blood or body fluids, use appropriate barrier protection such as disposable latex gloves, gowns, masks and eye coverings. Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant. Contact Time: Spray surface to be treated 3-4 seconds, from a distance of 6-8 inches, until surface is wet. Allow spray to dry without wiping or allow

For Use in Homes and Other Residential Dwellings, Hospitals, Nursing Homes, Clinics, Medical Facilities, Businesses, Hotels, Motels, Restaurants, Schools, Institutions and Commercial Establishments. Acts as a disinfectant on hard, non-porous environmental surfaces such as toilet seats, fixtures, glazed tile, wash bowls, shower stalls, walls, finished woodwork, counter tops, kitchen counter tops, tables and outside surfaces of appliances.

Features & Benefits

Bactericidal, *Virucidal, Fungicidal Cleans and Deodorizes Spray and Wipe Shining Clean

Use in nursery and sickroom, for garbage and diaper pails, for any washable hard, non-porous environmental surface that needs disinfecting as you clean.

The easy way to clean, disinfect and deodorize. Spray and wipe clean. Cleans and disinfects without scouring or scratching.

Country Fresh scent leaves the facility smelling great

Item Number & Unit of Measure

Item Number	HIL0103955
Unit of Measure	19 oz Can

Specifications

Color	White
Fragrance	Country Fresh

Germicidal Foaming Cleaner (continued)



spray to contact treated surface for a minimum of ten minutes prior to wiping. Disposal of Infectious Materials: Autoclave blood and other body fluids and dispose of according to federal, state and local regulations for infectious waste disposal. This product is not for use on medical devices or medical equipment surfaces. Efficacy: BACTERICIDAL ACTIVITY: 10 minute contact time, Staphylococcus aureus and Salmonella choleraesuis.