

QUESTION

1. A patient is admitted to the hospital with a diagnosis of acute myocardial infarction (AMI). The patient is in the intensive care unit (ICU) and is receiving intravenous (IV) therapy. The nurse is monitoring the patient's vital signs and the IV infusion. The patient's vital signs are stable, and the IV infusion is running at the prescribed rate. The nurse notices that the patient's oxygen saturation is 92% on 2L of oxygen. The nurse should:

1. Continue to monitor the patient's oxygen saturation.
2. Increase the oxygen flow to 4L.
3. Decrease the oxygen flow to 1L.
4. Notify the physician of the patient's oxygen saturation.

ANSWER

1. A patient is admitted to the hospital with a diagnosis of acute myocardial infarction (AMI). The patient is in the intensive care unit (ICU) and is receiving intravenous (IV) therapy. The nurse is monitoring the patient's vital signs and the IV infusion. The patient's vital signs are stable, and the IV infusion is running at the prescribed rate. The nurse notices that the patient's oxygen saturation is 92% on 2L of oxygen. The nurse should:

1. Continue to monitor the patient's oxygen saturation.
2. Increase the oxygen flow to 4L.
3. Decrease the oxygen flow to 1L.
4. Notify the physician of the patient's oxygen saturation.

QUESTION

1. A patient is admitted to the hospital with a diagnosis of acute myocardial infarction (AMI). The patient is in the intensive care unit (ICU) and is receiving intravenous (IV) therapy. The nurse is monitoring the patient's vital signs and the IV infusion. The patient's vital signs are stable, and the IV infusion is running at the prescribed rate. The nurse notices that the patient's oxygen saturation is 92% on 2L of oxygen. The nurse should:

1. Continue to monitor the patient's oxygen saturation.
2. Increase the oxygen flow to 4L.
3. Decrease the oxygen flow to 1L.
4. Notify the physician of the patient's oxygen saturation.

ANSWER

1. A patient is admitted to the hospital with a diagnosis of acute myocardial infarction (AMI). The patient is in the intensive care unit (ICU) and is receiving intravenous (IV) therapy. The nurse is monitoring the patient's vital signs and the IV infusion. The patient's vital signs are stable, and the IV infusion is running at the prescribed rate. The nurse notices that the patient's oxygen saturation is 92% on 2L of oxygen. The nurse should:

1. Continue to monitor the patient's oxygen saturation.
2. Increase the oxygen flow to 4L.
3. Decrease the oxygen flow to 1L.
4. Notify the physician of the patient's oxygen saturation.