



Item Number & Unit of Measure

Item Number	HIL0081989
Unit of Measure	1 fl oz Packet

Specifications

Color	Blue
Appearance	Clear
Fragrance	Lavender
Formula Type	Liquid
Dilution Ratio	1:64
PH	6.00 - 7.00
Non-Volatile Matter	7.50 - 8.50%
EPA Registration #	1839-169-1658

Safety

See safety datasheet (if applicable) and product label for safety information, handling and proper use.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not for use on critical and semi-critical medical device surfaces.

Surfaces which may contact food must be rinsed thoroughly with potable water after use.

GENERAL CLEANING - Apply Non-Acid Restroom Disinfectant/Cleaner to soiled area with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer. Non-Acid Restroom Disinfectant/Cleaner can be diluted at 2 ounces per gallon or applied at full strength. Thoroughly wet soiled surface and scrub as necessary. Non-Acid Restroom Disinfectant/Cleaner may be used to clean porous or non-porous surfaces. Non-Acid Restroom Disinfectant/Cleaner may be used in bathrooms, institutions, classrooms, hog farms, poultry and turkey farms or on tables, desks, chairs, garbage cans, shelves, sinks, shower doors and curtains, exercise equipment and wrestling mats. **DEODORIZATION** - To deodorize, apply Non-Acid Restroom Disinfectant/Cleaner as indicated under the heading **DISINFECTION**. **DISINFECTION** - To disinfect

Non-Acid restroom Disinfectant/Cleaner is a phosphate-free, safe for surfaces pH of 5.5-7.5 formulation designed to provide effective cleaning, deodorizing, and disinfection for hospitals, nursing homes, households, homes, schools and colleges, food processing plants, food service establishments, restaurants, transportation terminals, office buildings, manufacturing facilities, lodging establishments, hotels, retail business, and athletic/recreation facilities, sports stadiums, amphitheatres and convention centers where housekeeping is of prime importance in controlling cross-contamination from treated surfaces. Surfaces which may contact food must be rinsed thoroughly with potable water after use.

This product, when used as directed, is formulated to disinfect the following hard non-porous inanimate environmental surfaces: finished floors (sealed and unsealed floors), walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, vanity tops, shower stalls, bathtubs, and cabinets.

Non-Acid Restroom Disinfectant/Cleaner is effective against SARS-CoV-2 (COVID-19 virus) on hard non-porous surfaces in 1 minute.

Features & Benefits

EPA Registration No. 1839-169-1658

One-step, non-acid, germicidal detergent and deodorizer

Safe for surfaces pH of 5.5-7.5

Leaves a pleasant scent.

Intended for use with the Hillyard Arsenal One Dispensing systems.

Non-Acid Restroom Disinfectant/Cleaner (continued)



inanimate, hard non-porous surfaces add 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water. Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes or 1 minute for SARS-Related Coronavirus 2 (SARS-CoV-2), then remove excess liquid. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray mist. For visibly soiled areas, a pre-cleaning step is required. ***VIRUCIDAL ACTIVITY** - Non-Acid Restroom Disinfectant/Cleaner when used on environmental, inanimate, hard, non-porous surfaces exhibits effective virucidal activity against HIV-1, HIV-2, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Type 1 (causative agent of fever blisters), Herpes Simplex Type 2 (genital), Influenza A2/Hong Kong, Vaccinia, Rotavirus, Human Coronavirus (ATCC VR-740, Strain 229E), SARS Associated Coronavirus, Rabies Virus, Paramyxovirus (Mumps), Bovine Viral Diarrhea Virus (BVDV), Pseudorabies, Bovine Rhinotracheitis, Feline Leukemia, Feline Picornavirus, Canine Distemper Virus, Avian Influenza A Virus, Porcine Respiratory & Reproductive Syndrome Virus (PRRSV), and SARS-Related Coronavirus 2 (SARS-CoV-2) (USA-WA1/2020) (BEI NR-52281). Add 2 ounces of the product per gallon of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a cloth, mop, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow the surface to remain visibly wet for 10 minutes, then remove excess liquid. **KILLS HIV-1, HIV-2, HBV, and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human

Immunodeficiency Virus Type 1 or Type 2 (HIV-1 or HIV-2) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HIV-2, HBV, and HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of Non-Acid Restroom Disinfectant/Cleaner. **CONTACT TIME:** Allow surface to remain visibly wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal. **FUNGICIDAL ACTIVITY** - At the 2 ounce per gallon dilution Non-Acid Restroom Disinfectant/Cleaner is fungicidal against the pathogenic fungi, Trichophyton interdigitale, formerly known as Trichophyton mentagrophytes (Athlete's Foot Fungus) (a cause of ringworm of the foot) and Candida albicans. Apply solution with a cloth, sponge or hand pump trigger sprayer to hard, non-porous surfaces found in bathrooms, shower stalls, locker rooms, exercise facilities or other clean, hard non-porous surfaces commonly contacted by bare feet. Allow the surface to remain visibly wet for 10 minutes, then remove excess liquid. Diluted product should be applied daily or more frequently with heavy facility use. **MILDEWSTAT** - To control mold and mildew (Aspergillus niger) and the odors they cause on pre-cleaned, hard, non-porous inanimate surfaces add 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water. Apply solution with a cloth, mop, sponge or hand pump trigger sprayer making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth appears. **BACTERICIDAL ACTIVITY** - At the 2 ounce per gallon dilution Non-

Non-Acid Restroom Disinfectant/Cleaner (continued)



Acid Restroom Disinfectant/Cleaner demonstrates effective disinfectant activity against the organisms: *Pseudomonas aeruginosa* PRD-10, *Salmonella (choleraesuis) enterica*, *Staphylococcus aureus*, *Staphylococcus aureus* (clinical isolate), *Bordetella bronchiseptica*, *Corynebacterium ammoniagenes*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter cloacae* (clinical isolate), *Enterococcus faecalis*, *Enterococcus faecalis* (clinical isolate), *Escherichia coli*, *Escherichia coli* (clinical isolate), *Fusobacterium necrophorum*, *Klebsiella pneumoniae* subsp. *pneumoniae*, *Lactobacillus casei* subsp. *raamnosus*, *Listeria monocytogenes*, *Pasteurella multocida*, *Proteus vulgaris*, *Proteus mirabilis* (ATCC 9921), *Proteus mirabilis* (ATCC 25933), *Salmonella enterica* subsp. *enterica* serotype *paratyphi B*, *Salmonella enterica* subsp. *enterica* serotype *typhi*, *Salmonella enterica* subsp. *enterica* serotype *typhimurium*, *Salmonella enterica* subsp. *enterica* serotype *pullorum*, *Serratia marcescens*, *Shigella sonnei*, *Shigella flexneri* Type 2b, *Shigella dysenteriae*, *Staphylococcus aureus* subsp. *aureus*, *Staphylococcus epidermidis*, *Staphylococcus epidermidis* (clinical isolate), *Streptococcus pyogenes* (Clinical - Flesh Eating Strain BIRD M3), *Streptococcus pyogenes* Group A, *Xanthomonas maltophilia* (clinical isolate), Vancomycin resistant *Enterococcus faecalis* (VRE), Methicillin resistant *Staphylococcus aureus* (MRSA), Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) (NRS 123 Genotype USA400), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) (NRS 384 Genotype USA300). **BACTERICIDAL STABILITY OF USE-SOLUTION:** Tests confirm that Non-Acid Restroom Disinfectant/Cleaner when diluted in hard water up to 400 ppm (as CaCO₃) remains effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella (choleraesuis) enterica* for up to 30 days when stored in a sealed container such as a

spray bottle at room temperature. If Non-Acid Restroom Disinfectant/Cleaner becomes visibly dirty or contaminated, the use-solution must be discarded and fresh use-solution prepared. Always use clean, properly labeled containers when diluting Non-Acid Restroom Disinfectant/Cleaner. Bactericidal stability of the use-solution does not apply to open containers such as buckets or pails. **To disinfect toilet bowls** - Remove visible filth or soils from surfaces with bowl brush. Add 2 ounces of Non-Acid Restroom Disinfectant/Cleaner to the bowl water. Brush or swab the bowl completely using a scrub brush or toilet mop making sure to get under the rim. Let stand for 10 minutes and flush. **To disinfect food processing premises:** Before using Non-Acid Restroom Disinfectant/Cleaner, food products and packaging materials must be removed from the area or carefully protected. For floors, walls, and storage areas, add 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes, then remove excess liquid. After use, all surfaces in the area must be thoroughly rinsed with potable water. **To disinfect restaurant (food service establishment) food contact surfaces:** countertops, appliances, and tables, add 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow the surface to remain visibly wet for 10 minutes, then remove excess liquid and rinse the surface with potable water. Non-Acid Restroom Disinfectant/Cleaner cannot be used to clean the following food contact surfaces: utensils, glassware and dishes. **POULTRY AND SWINE PREMISE DISINFECTION Site Preparation:** The first step in any on-going disinfection program must be the removal of visible contamination and debris. This is

accomplished by using a shovel, broom, or vacuum, depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of visible organic matter. Once the visible debris is eliminated, thoroughly clean all surfaces with soap or detergent and rinse with water.

Cleaning And Disinfection: For all general cleaning and disinfection, use 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water. Apply Non-Acid Restroom Disinfectant/Cleaner using a cloth, mop, or low pressure coarse sprayer so as to thoroughly wet surface to be disinfected. Allow to remain visibly wet for 10 minutes and then let air dry. For visibly soiled areas, a pre-cleaning step is required. Prepare a fresh solution after each use.

APPLICATIONS AND USE DILUTION

Poultry And Swine Premise: Remove all animals and feeds from premises, trucks, cars, coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water. Saturate surfaces with the disinfecting solution for a period of 10 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

Hatcheries: Use 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water to treat the following hard non-porous surfaces: hatching, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more. Rinse with potable

water before reuse. Then allow to air dry.

Vehicles: Clean all vehicles including mats, crates, cabs, and wheels with high pressure water and Non-Acid Restroom Disinfectant/Cleaner. Use 2 ounces per gallon to treat all vehicles. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more.

Veterinary Practice/Animal Care/Animal

Laboratory/Zoos/Pet Shop/Kennels Disinfection
Directions: For cleaning and disinfecting the following hard non-porous surfaces: equipment not used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feed from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces occupied or traversed by animals.

Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

Disinfection of Barber/Salon Tools

Directions: Immerse pre-cleaned barber/salon tools (combs, brushes, razors, manicure/pedicure tools, clippers, scissors, trimmer blades) in a use-solution (or equivalent dilution) of 2 ounces of Non-Acid Restroom Disinfectant/Cleaner per gallon of water for at least 10 minutes. Rinse instruments thoroughly and dry before reuse. A fresh use-solution must be prepared daily or more often if the use-solution becomes visibly dirty. **NOTE:**

Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed from the use-solution after 10 minutes, rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged immersion may cause damage to stainless steel or metal instruments.

To Control Mold and Mildew on Large, Inflatable, Non-Porous Plastic and Rubber Structures (animals, promotional items,

Non-Acid Restroom Disinfectant/Cleaner (continued)



moonwalk, slides, obstacle course play equipment, exercise equipment, and wrestling mats): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 8 ounces of Non-Acid Restroom Disinfectant/Cleaner per 4 gallons of water (2 ounces per gallon of water) (850 ppm active quaternary) (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

Bovine Viral Diarrhea Virus (BVDV), Pseudorabies, Bovine Rhinotracheitis, Feline Leukemia, Feline Picornavirus, Canine Distemper Virus, Avian Influenza A Virus, Porcine Respiratory & Reproductive Syndrome Virus (PRRSV), and 1-minute contact time: SARS-Related Coronavirus 2 (SARS-CoV-2) (USA-WA1/2020) (BEINR-52281).

HMIS Ratings

Flammability RTU	0
Health RTU	1
Reactivity RTU	0

Efficacy

BACTERICIDAL ACTIVITY: 10-minute contact time, *Pseudomonas aeruginosa* PRD-10, *Salmonella* (choleraesuis) enterica, *Staphylococcus aureus*, *Staphylococcus aureus* (clinical isolate), *Bordetella bronchiseptica*, *Corynebacterium ammoniagenes*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter cloacae* (clinical isolate), *Enterococcus faecalis*, *Enterococcus faecalis* (clinical isolate), *Escherichia coli*, *Escherichia coli* (clinical isolate), *Fusobacterium necrophorum*, *Klebsiella pneumoniae* subsp. pneumoniae, *Lactobacillus casei* subsp. rhamnosus, *Listeria monocytogenes*, *Pasteurella multocida*, *Proteus vulgaris*, *Proteus mirabilis* ATCC9921, *Proteus mirabilis* ATCC 25933, *Salmonella* (paratyphi B) enterica, *Salmonella* (typhi) enterica, *Salmonella* (typhimurium) enterica, *Salmonella* (pullorum) enterica, *Serratia marcescens*, *Shigella sonnei*, *Shigella flexneri* Type 2b, *Shigella dysenteriae*, *Staphylococcus aureus* subsp. aureus, *Staphylococcus epidermidis*, *Staphylococcus epidermidis* (clinical isolate), *Streptococcus pyogenes* (Clinical - Flesh Eating Strain BIRD M3), *Streptococcus pyogenes* Group A, *Xanthomonas maltophilia* (clinical isolate), Vancomycin resistant *Enterococcus faecalis* (VRE), Methicillin resistant *Staphylococcus aureus* (MRSA), Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) NRS123 Genotype USA400, Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) NRS 384 Genotype USA300. **FUNGICIDAL ACTIVITY: 10-minute** contact time, *Trichophyton mentagrophytes* (Athlete's Foot Fungus) and *Candida albicans*. **MILDEWSTAT: 10-minute** contact time: *Aspergillus niger*. **VIRUCIDAL ACTIVITY: 10-minute** contact time: HIV-1, HIV-2, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Type 1 (causative agent of fever blisters), Herpes Simplex Type 2 (genital), Influenza A2/Hong Kong, Vaccinia, Rotavirus, Human Coronavirus (ATCC VR-740, Strain 229E), SARS Associated Coronavirus, Rabies Virus, Paramyxovirus (Mumps),