

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a 10% weight loss, muscle wasting, and a 10% decrease in albumin. The patient has a 10% decrease in albumin. The patient has a 10% decrease in albumin.

Parameter	Normal	Abnormal
Albumin	3.5-5.0 g/dL	2.5-3.0 g/dL
Weight	70 kg	63 kg
Albuminuria	<30 mg/day	>30 mg/day
Proteinuria	<300 mg/day	>300 mg/day
Proteinuria	<300 mg/day	>300 mg/day
Proteinuria	<300 mg/day	>300 mg/day

2. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a 10% weight loss, muscle wasting, and a 10% decrease in albumin. The patient has a 10% decrease in albumin. The patient has a 10% decrease in albumin.

ANSWER



3. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a 10% weight loss, muscle wasting, and a 10% decrease in albumin. The patient has a 10% decrease in albumin. The patient has a 10% decrease in albumin.

4. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a 10% weight loss, muscle wasting, and a 10% decrease in albumin. The patient has a 10% decrease in albumin. The patient has a 10% decrease in albumin.