

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and a 10-degree increase in serum bilirubin. The patient has a history of chronic liver disease and is currently on a low-fat diet. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

| Test | Result |
|----------------------|------------|
| Albumin | 2.5 g/dL |
| Bilirubin | 1.5 mg/dL |
| ALT | 100 U/L |
| AST | 150 U/L |
| Alkaline phosphatase | 100 U/L |
| Gamma-GT | 150 U/L |
| Prothrombin time | 15 seconds |
| INR | 1.5 |
| PTAP | 1.5 |
| PTAP/APTT ratio | 1.5 |
| PTAP/APTT ratio | 1.5 |

2. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and a 10-degree increase in serum bilirubin. The patient has a history of chronic liver disease and is currently on a low-fat diet. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

| Test | Result |
|----------------------|------------|
| Albumin | 2.5 g/dL |
| Bilirubin | 1.5 mg/dL |
| ALT | 100 U/L |
| AST | 150 U/L |
| Alkaline phosphatase | 100 U/L |
| Gamma-GT | 150 U/L |
| Prothrombin time | 15 seconds |
| INR | 1.5 |
| PTAP | 1.5 |
| PTAP/APTT ratio | 1.5 |
| PTAP/APTT ratio | 1.5 |

3. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and a 10-degree increase in serum bilirubin. The patient has a history of chronic liver disease and is currently on a low-fat diet. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

| Test | Result |
|----------------------|------------|
| Albumin | 2.5 g/dL |
| Bilirubin | 1.5 mg/dL |
| ALT | 100 U/L |
| AST | 150 U/L |
| Alkaline phosphatase | 100 U/L |
| Gamma-GT | 150 U/L |
| Prothrombin time | 15 seconds |
| INR | 1.5 |
| PTAP | 1.5 |
| PTAP/APTT ratio | 1.5 |
| PTAP/APTT ratio | 1.5 |

ANSWER

1. The patient's laboratory studies are consistent with alcoholic liver disease. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

| Test | Result |
|----------------------|------------|
| Albumin | 2.5 g/dL |
| Bilirubin | 1.5 mg/dL |
| ALT | 100 U/L |
| AST | 150 U/L |
| Alkaline phosphatase | 100 U/L |
| Gamma-GT | 150 U/L |
| Prothrombin time | 15 seconds |
| INR | 1.5 |
| PTAP | 1.5 |
| PTAP/APTT ratio | 1.5 |
| PTAP/APTT ratio | 1.5 |

2. The patient's laboratory studies are consistent with alcoholic liver disease. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

| Test | Result |
|----------------------|------------|
| Albumin | 2.5 g/dL |
| Bilirubin | 1.5 mg/dL |
| ALT | 100 U/L |
| AST | 150 U/L |
| Alkaline phosphatase | 100 U/L |
| Gamma-GT | 150 U/L |
| Prothrombin time | 15 seconds |
| INR | 1.5 |
| PTAP | 1.5 |
| PTAP/APTT ratio | 1.5 |
| PTAP/APTT ratio | 1.5 |

3. The patient's laboratory studies are consistent with alcoholic liver disease. The patient's physical examination is unremarkable. The patient's laboratory studies are as follows:

| Test | Result |
|----------------------|------------|
| Albumin | 2.5 g/dL |
| Bilirubin | 1.5 mg/dL |
| ALT | 100 U/L |
| AST | 150 U/L |
| Alkaline phosphatase | 100 U/L |
| Gamma-GT | 150 U/L |
| Prothrombin time | 15 seconds |
| INR | 1.5 |
| PTAP | 1.5 |
| PTAP/APTT ratio | 1.5 |
| PTAP/APTT ratio | 1.5 |