



**Code:** 9676 **Size:** 4.55 oz (129 g) Powder

# **Myo-Inositol Plus**

## with DC-Inositol, Folate, and B12

# Supports Ovarian and Reproductive System Function\*

- Provides recommended 40:1 ratio of myo-inositol to D-chiro-inositol
- Contains naturally occurring forms of inositol
- Includes methylcobalamin and activated folate (L-5-MTHF) for healthy fertility and pregnancy\*
- Adequate folate in healthful diets may reduce a woman's risk of having a child with a brain or spinal cord birth defect
- Unflavored powder allows for easy twice-per-day dosing

## PRODUCT SUMMARY

Myo-Inositol Plus provides the two most common, naturally occurring, and complementary forms of inositol: myoinositol and D-chiro-inositol. Both forms of inositol have been shown to support ovarian, metabolic, and endocrine function in women.\* Myo-inositol and D-chiro-inositol have been shown to support both ovarian function, including cycle regularity and normal LH levels, and metabolic function, including HOMA index and insulin.\* Additionally, D-chiro-inositol supports endocrine function such as normal androgen levels, as well as blood pressure already within the normal range.\*<sup>1.9</sup>

In Myo-Inositiol Plus, these complementary nutrients are combined in the 40:1 ratio that has been recommended by the International Consensus Conference on myo-inositol and D-chiro-inositol.<sup>10</sup>

This product also contains methylcobalamin and activated folate (L-5-MTHF) for healthy fertility and pregnancy,\* both of which are active forms that do not require metabolism for use. Adequate folate in healthful diets may reduce a woman's risk of having a child with a brain or spinal cord birth defect.



**To Place Your Order Email:** customersupport@bioclinicnaturals.com **Call:** 1.877.433.9860 • **Fax:** 1.877.433.9862







Supplement Facts		
Serving Size 1 Scoop (2.15 g) Servings Per Container 60		
Amo	ount Per Serving	% Daily Value
Folate (L-5-Methyltetrahydrofolat Calcium Salt) (20	e, 340 mcg DFE 10 mcg Folic Acid)	85%
Vitamin B12 (Methylcobalamin)	1.5 mcg	63%
Myo-Inositol	2 g	**
D-(+)-Chiro Inositol	50 mg	**
** Daily Value not established.		

Other ingredients: Organic rice hull concentrate.

Suggested Usage: Stir 1 scoop (2.15 g) into 250 ml of water 1-2 times per day or as directed by a health care professional. Myo-Inositol Plus has been formulated to support ovarian and reproductive system function,\* and to aid in promoting healthy glucose metabolism already within the normal range.\*

Caution: Consult your health care professional prior to use if you are pregnant, trying to become pregnant, breastfeeding, taking medication, have a medical condition, or anticipate surgery. Keep out of reach of children.

Drug Interactions: Myo-inositol has been shown to enhance the efficacy of clomiphene. When given with folic acid, it improved ovulation rates (65.5%) compared to clomiphene alone (42%), as well as pregnancy rates (53.8% combined use vs. 42.2% clomiphene alone).<sup>11</sup> Medication dosage to control hyperglycemia may need monitoring, as myo-inositol has benefits on insulin sensitivity similar to metformin.<sup>12</sup>

Contains no artificial colors, preservatives, or sweeteners; no dairy, starch, sugar, wheat, gluten, yeast, soy, egg, fish, shellfish, animal products, salt, tree nuts, or GMOs. Suitable for vegetarians/vegans. Sealed for your protection. Do not use if seal is broken. For freshness, store in a cool, dry place.

### References

- Gerli, S., Papaleo, E., Ferrari, A., et al. (2007). Eur Rev Med Pharmacol Sci, 11(5), 347-354.
- Unfer, V., Facchinetti, F., Orrù, B., et al. (2017). Endocr Connect, 6(8), 647-658. 3
- Costantino, D., Minozzi, G., Minozzi, E., et al. (2009). Eur Rev Med Pharmacol Sci, 13(2), 105-110.
- Nordio, M., & Proietti, E. (2012). Eur Rev Med Pharmacol Sci, 16(5), 575-581.
- 5 Colazingari, S., Treglia, M., Najjar, R., et al. (2013). Arch Gynecol Obstet, 288(6), 1405-1411.
- Pizzo, A., Laganà, A.S., & Barbaro, L. (2014). *Gynecol Endocrinol, 30*(3), 205-208. Formuso, C., Stracquadanio, M., & Ciotta, L. (2015). *Minerva Ginecol, 67*(4), 321-325 6.
- 8 Laganà, A.S., Barbaro, L., & Pizzo, A. (2015). Arch Gynecol Obstet, 291(5), 1181-1186.
- Genazzani, A.D., Santagni, S., Rattighieri, E., et al. (2014). Gynecol Endocrinol, 30(6), 9. 438-443
- 10. Facchinetti, F., Bizzarri, M., Benvenga, S., et al. (2015). Eur J Obstet Gynecol Reprod Biol, 195. 72-76.
- 11. Rolland, A.L., Peigné, M., Plouvier, P., et al. (2017). Eur Rev Med Pharmacol Sci, 21(2 Suppl), 10-14
- 12. Fruzzetti, E. Perini, D., Russo, M., et al. (2017), Gvnecol Endocrinol, 33(1), 39-42.



#### · GUARANTEED ·

Bioclinic Naturals® products are guaranteed to meet or exceed Good Manufacturing Practices (GMP) of the U.S. Food and Drug Administration (FDA), Health Canada, and the Therapeutic Goods Administration (TGA) of Australia.



PRODUCT OF CANADA Manufactured for and distributed by Bioclinic Naturals® U S 14224 167th Ave. SE, Monroe WA, USA 98272 bioclinicnaturals.com

FOR PROFESSIONAL USE ONLY. \*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease. © All Rights Reserved Bioclinic Naturals® 2020. July 21, 2020. 9224506

Head office Assured Natural Distribution Inc., 104 - 3686 Bonneville Place, Burnaby, BC, Canada V3N 4T6 | U.S. Distribution office 14224 167th Avenue SE, Monroe, WA, USA 98272 Customer service 1.877.433.9860 · Fax 1.877.433.9862 · Email customersupport@bioclinicnaturals.com