

BYNACOLIN[®] B12

100 mg Methylcobalamin
per 500 mg Tablet



Directions:
Take 1 tablet orally, once daily with food. Do not crush or chew the tablet. If you are unable to swallow the tablet whole, you may crush the tablet and mix it with a small amount of soft food (e.g., applesauce) and swallow the mixture.

Warnings:
Do not use if you are allergic to any of the ingredients listed in the inactive ingredients section. Do not use if you are pregnant or breastfeeding, or if you are taking other medications that may interact with Bynacolin B12.

Side Effects:
Common side effects include headache, dizziness, and nausea. If you experience any of these symptoms, contact your healthcare provider.

Other Information:
This product is a prescription drug. It is intended for use in the United States only. Do not use if the seal is broken or if the tablets are discolored.

Ingredients:
Active ingredient: Methylcobalamin (100 mg)
Inactive ingredients: Lactose monohydrate, croscarmellose sodium, hydroxypropyl methylcellulose, polyethylene glycol 400, and hydroxypropyl cellulose.

How to Use:
See the directions for use on the label.

Storage:
Store at room temperature (20° to 25°C).

Keep out of reach of children.

See important information about this medicine on the label.

© 2023 BSN. All rights reserved.

BYNACOLIN[®] B12 is a registered trademark of BSN.

100 mg Methylcobalamin per 500 mg Tablet

Tablets, 500 mg

Net Content: 30 Tablets

U.S. Patent Nos. 8,912,118; 8,912,119; 8,912,120