

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, anorexia, and weakness. Physical examination shows a 10% weight loss, tachycardia, and a 30-degree angle of trunk extension. Laboratory studies show a total bilirubin of 2.5 mg/dL, a prothrombin time of 18 seconds, and a serum albumin of 2.5 g/dL. The patient is most likely suffering from:

A. Acute liver failure
B. Chronic liver disease
C. Hemolytic anemia
D. Iron deficiency anemia

2. A 65-year-old man with a long history of alcohol abuse presents with a 2-week history of weight loss, anorexia, and weakness. Physical examination shows a 10% weight loss, tachycardia, and a 30-degree angle of trunk extension. Laboratory studies show a total bilirubin of 2.5 mg/dL, a prothrombin time of 18 seconds, and a serum albumin of 2.5 g/dL. The patient is most likely suffering from:

A. Acute liver failure
B. Chronic liver disease
C. Hemolytic anemia
D. Iron deficiency anemia

ANSWER: B

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, anorexia, and weakness. Physical examination shows a 10% weight loss, tachycardia, and a 30-degree angle of trunk extension. Laboratory studies show a total bilirubin of 2.5 mg/dL, a prothrombin time of 18 seconds, and a serum albumin of 2.5 g/dL. The patient is most likely suffering from:

A. Acute liver failure
B. Chronic liver disease
C. Hemolytic anemia
D. Iron deficiency anemia

2. A 65-year-old man with a long history of alcohol abuse presents with a 2-week history of weight loss, anorexia, and weakness. Physical examination shows a 10% weight loss, tachycardia, and a 30-degree angle of trunk extension. Laboratory studies show a total bilirubin of 2.5 mg/dL, a prothrombin time of 18 seconds, and a serum albumin of 2.5 g/dL. The patient is most likely suffering from:

A. Acute liver failure
B. Chronic liver disease
C. Hemolytic anemia
D. Iron deficiency anemia

ANSWER: B