

QUESTION
A 55-year-old male with a long history of hypertension and hyperlipidemia presents with a 3-month history of progressive weakness and fatigue. He reports a 15-pound weight loss over the past 6 months. Physical examination reveals tachycardia, a bounding pulse, and a systolic blood pressure of 160/90 mmHg. Laboratory studies show a hemoglobin of 12 g/dL, a hematocrit of 38%, and a serum ferritin of 100 ng/mL. The patient's thyroid function tests are within normal limits. What is the most likely diagnosis?

ANSWER
Aortic regurgitation

QUESTION
A 68-year-old male with a long history of hypertension and hyperlipidemia presents with a 3-month history of progressive weakness and fatigue. He reports a 15-pound weight loss over the past 6 months. Physical examination reveals tachycardia, a bounding pulse, and a systolic blood pressure of 160/90 mmHg. Laboratory studies show a hemoglobin of 12 g/dL, a hematocrit of 38%, and a serum ferritin of 100 ng/mL. The patient's thyroid function tests are within normal limits. What is the most likely diagnosis?

ANSWER
Aortic regurgitation

ANSWERS

QUESTION
A 55-year-old male with a long history of hypertension and hyperlipidemia presents with a 3-month history of progressive weakness and fatigue. He reports a 15-pound weight loss over the past 6 months. Physical examination reveals tachycardia, a bounding pulse, and a systolic blood pressure of 160/90 mmHg. Laboratory studies show a hemoglobin of 12 g/dL, a hematocrit of 38%, and a serum ferritin of 100 ng/mL. The patient's thyroid function tests are within normal limits. What is the most likely diagnosis?

ANSWER
Aortic regurgitation