



LYSOL® IC™

Quaternary Disinfectant Cleaner

Virucidal, fungicidal and bactericidal, this multi-purpose, hospital disinfectant cleans and disinfects hard, nonporous surfaces. The economical, concentrated formula makes up to 256 gallons (1:256). Use on floors, walls, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, etc.) and other hard, nonporous surfaces.

Features & Benefits

- Controls Mold & Mildew
- Kills E.Coli, Salmonella, Listeria, MRSA, Staph and Strep.*
- Use in healthcare facilities, locker rooms, restaurants, schools, hotels



* When used as directed.





LYSOL® IC™ Quaternary Disinfectant Cleaner

Use Instructions

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is not for use on medical device surfaces.

DISINFECTION/CLEANING/DEODORIZING DIRECTIONS:

- Remove heavy soil deposits from surface
- Then thoroughly wet surface with a use-solution of 1/2 ounce of concentrate per gallon of water or equivalent. The use-solution can be applied with a cloth, mop, sponge, or coarse spray, or soaking.
- For sprayer applications, use a coarse spray device
- Spray 6-8 inches from the surface, rub with a brush, cloth or sponge. Do not breathe spray. Let solution remain on surface for a minimum of 10 minutes. Rinse or allow to air dry.
- Rinsing of floors is not necessary unless they are to be waxed or polished. Food contact surfaces must be thoroughly rinsed with potable water. This product must not be used to clean the following food contact surfaces: utensils, glassware and dishes.
- For Human Coronavirus, treated surfaces must remain wet for 1 minute. Wipe dry with a clean cloth, sponge or mop or allow to air dry.
- For Influenza Virus Type A, treated surfaces must remain wet for 2 minutes. Wipe dry with a clean cloth, sponge or mop or allow to air dry.
- Prepare a fresh solution daily or more often if the solution becomes visibly dirty or diluted
- *KILLS HIV-1 (AIDS VIRUS) AND HBV (HEPATITIS B VIRUS) AND HCV (HEPATITIS C VIRUS) ON PRECLEANED, HARD NON-POROUS ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR DISINFECTION, CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OR HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

- PERSONAL PROTECTION: Disposable protective gloves, gowns, face masks and eye coverings must be worn during all cleaning of blood/body fluids and during decontamination procedures
- CLEANING PROCEDURES: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of disinfectant
- CONTACT TIME: Mix 1/2 oz. of this product per gallon of water (or equivalent dilution). HIV-1 (AIDS virus) is inactivated after a contact time of 4 minutes at 25° C (77° F). HBV and HCV are inactivated after a 10 minute contact time. Use a 10 minute contact time for all other viruses, fungi, and bacteria listed.

DISPOSAL OF INFECTIOUS MATERIALS:

- Blood/body fluids must be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal
- TOILET BOWLS: Swab bowl with brush to remove heavy soil prior to cleaning or disinfecting. Clean by applying use-solution around the bowl and up under the rim. Stubborn stains may require brushing. To disinfect, first pre-clean to remove heavy soil, then remove or expel over the inner trap the residual bowl water. Pour in 3 oz. of the use-solution. Swab the bowl completely using a scrub brush or mop, making sure to get under rim. Let stand for 10 minutes or overnight, then flush.
- FUNGICIDAL DIRECTIONS: For use in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities follow disinfection directions.
- MILDEWSTATIC INSTRUCTIONS: Will effectively control the growth of mold and mildew plus the odors caused by them when applied to hard, nonporous surfaces such as walls, floors, and table tops. Apply use-solution of 1/2 oz. per gallon of water with a cloth, mop, sponge, or coarse spray. Make sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS DISINFECTION DIRECTIONS:

- For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc.
- Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls, and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances.
- Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 1/2 oz. of LYSOL® Brand I.C.™ Quaternary Disinfectant Cleaner per gallon of water (or equivalent dilution) for a period of 10 minutes. Wipe or allow to air dry. Ventilate buildings and other closed spaces.
- Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

FOR HEAVY DUTY CLEANING AND/OR ODOR CONTROL:

- When greater cleaning is desired, use 1 - 2 oz. of LYSOL® Brand I.C.™ Quaternary Disinfectant Cleaner per gallon of water. Heavily soiled areas may require repeated cleaning before treatment.

CLEANING AND DISINFECTING HARD NONPOROUS SURFACES ON PERSONAL PROTECTIVE EQUIPMENT (RESPIRATORS):

- Pre-clean equipment if heavily soiled to ensure proper surface contact. Add 1/2 oz. of this product to one gallon of water. Gently mix for a uniform solution. Apply solution to hard, nonporous surfaces of the respirator with a brush, coarse spray device, sponge or by immersion. Thoroughly wet all surfaces to be disinfected. Treated surfaces must remain wet for 10 minutes. Remove excess solution from equipment prior to storage. Comply with all OSHA regulations for cleaning respiratory protection equipment (29 CFR 1910.134)

Technical data	LYSOL® IC™ Quaternary Disinfectant Cleaner
Color/Form	Amber, Liquid
pH	7.2 to 8.2
Shelf Life	2 Years

Product	Pack size	Dilution	Product code
LYSOL® IC™ Quaternary Disinfectant Cleaner	4 x 1 gal. Containers	1:256	CB749832

Safe handling

Please make sure your employees read and understand the product label and Safety Data Sheet before using this product. The label contains directions for use; and both the label and SDS contain hazard warnings, precautionary statements and first aid procedures. SDS are available online at www.diversey.com or by calling 888.352.2249. Improper use or dilution may result in damage to surfaces and may result in health and physical hazards that match those of the concentrate. Please refer to the Diversey HazMat Library, only available through Internet Explorer, <http://naextranet.diversey.com/dot/>, for up-to-date shipping information.