

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a bilateral lower-extremity edema, a bilateral lower-extremity pitting edema, and a bilateral lower-extremity pitting edema. The patient's laboratory studies are as follows:

Test	Result
White blood cell count	12,000/mm ³
Hemoglobin	10 g/dL
Hematocrit	30%
Serum albumin	2.5 g/dL
Serum total protein	5.5 g/dL
Serum bilirubin	2.5 mg/dL
Serum aspartate aminotransferase (AST)	100 U/L
Serum alanine aminotransferase (ALT)	150 U/L
Serum creatinine	1.5 mg/dL
Serum prothrombin time (PT)	15 seconds
Serum international normalized ratio (INR)	1.5
Serum ammonia	100 μmol/L
Serum lactate dehydrogenase (LDH)	1000 U/L
Serum ferritin	1000 ng/mL
Serum ferritin	1000 ng/mL
Serum ferritin	1000 ng/mL

2. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a bilateral lower-extremity edema, a bilateral lower-extremity pitting edema, and a bilateral lower-extremity pitting edema. The patient's laboratory studies are as follows:

Test	Result
White blood cell count	12,000/mm ³
Hemoglobin	10 g/dL
Hematocrit	30%
Serum albumin	2.5 g/dL
Serum total protein	5.5 g/dL
Serum bilirubin	2.5 mg/dL
Serum aspartate aminotransferase (AST)	100 U/L
Serum alanine aminotransferase (ALT)	150 U/L
Serum creatinine	1.5 mg/dL
Serum prothrombin time (PT)	15 seconds
Serum international normalized ratio (INR)	1.5
Serum ammonia	100 μmol/L
Serum lactate dehydrogenase (LDH)	1000 U/L
Serum ferritin	1000 ng/mL
Serum ferritin	1000 ng/mL
Serum ferritin	1000 ng/mL

3. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a bilateral lower-extremity edema, a bilateral lower-extremity pitting edema, and a bilateral lower-extremity pitting edema. The patient's laboratory studies are as follows:

ANSWERS



4. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, weakness, and confusion. On physical examination, there is a bilateral lower-extremity edema, a bilateral lower-extremity pitting edema, and a bilateral lower-extremity pitting edema. The patient's laboratory studies are as follows: