



**Avert®**

**Sporicidal Disinfectant Cleaner**

<b>ACTIVE INGREDIENT:</b>	
Sodium hypochlorite .....	1.312%
<b>OTHER INGREDIENTS</b> .....	98.688%
<b>TOTAL</b> .....	100.000%

1.25% available Chlorine

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

See additional precautionary statements on back panel.

SEE ORIGINAL CONTAINER FOR COMPLETE DIRECTIONS FOR USE AND ADDITIONAL INFORMATION

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR FURTHER TREATMENT ADVICE.**  
1-800-851-7145

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOT FOR RESALE

**DIRECTIONS FOR USE:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Special Instructions for Cleaning Prior to Disinfection against Clostridium difficile spores**

**Personal Protection:** Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

**Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the sporicidal product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

**Infectious Materials Disposal:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

**Disinfection Procedure:** Apply product by coarse trigger sprayer to hard, nonporous inanimate surfaces. Spray 6 – 8 inches from surface, making sure to wet surfaces thoroughly. Surfaces must remain wet for 4 minutes. Wipe surfaces and let air dry.

**STORAGE AND DISPOSAL**

Do not contaminate food or feed by storage, disposal, or cleaning of equipment.

**STORAGE:** Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Wrap empty container and put in trash or offer for recycling if available.

**KEEP OUT OF REACH OF CHILDREN FOR INSTITUTIONAL USE**

**ALWAYS RINSE BOTTLE AND AIR DRY THOROUGHLY BEFORE REFILLING THIS BOTTLE SHOULD NOT BE USED WITH ANY OTHER PRODUCT.**

**CONTENTS OF THIS CONTAINER PREPARED BY USER.**

EPA Reg. No. 70627-72



8 07174 53060 5

©2018 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

LBL91338 (18/113) PRINTED IN THE U.S.A.



**Avert®**

**Sporicidal Disinfectant Cleaner**

<b>ACTIVE INGREDIENT:</b>	
Sodium hypochlorite .....	1.312%
<b>OTHER INGREDIENTS</b> .....	98.688%
<b>TOTAL</b> .....	100.000%

1.25% available Chlorine

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

See additional precautionary statements on back panel.

SEE ORIGINAL CONTAINER FOR COMPLETE DIRECTIONS FOR USE AND ADDITIONAL INFORMATION

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR FURTHER TREATMENT ADVICE.**  
1-800-851-7145

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOT FOR RESALE

**DIRECTIONS FOR USE:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Special Instructions for Cleaning Prior to Disinfection against Clostridium difficile spores**

**Personal Protection:** Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

**Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the sporicidal product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

**Infectious Materials Disposal:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

**Disinfection Procedure:** Apply product by coarse trigger sprayer to hard, nonporous inanimate surfaces. Spray 6 – 8 inches from surface, making sure to wet surfaces thoroughly. Surfaces must remain wet for 4 minutes. Wipe surfaces and let air dry.

**STORAGE AND DISPOSAL**

Do not contaminate food or feed by storage, disposal, or cleaning of equipment.

**STORAGE:** Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Wrap empty container and put in trash or offer for recycling if available.

**KEEP OUT OF REACH OF CHILDREN FOR INSTITUTIONAL USE**

**ALWAYS RINSE BOTTLE AND AIR DRY THOROUGHLY BEFORE REFILLING THIS BOTTLE SHOULD NOT BE USED WITH ANY OTHER PRODUCT.**

**CONTENTS OF THIS CONTAINER PREPARED BY USER.**

EPA Reg. No. 70627-72



8 07174 53060 5

©2018 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

LBL91338 (18/113) PRINTED IN THE U.S.A.