

QUESTION

1. A patient with a long history of alcohol abuse presents with a 2-week history of weight loss, anorexia, and weakness. The patient has a 30-year history of alcohol abuse and has been drinking approximately 100 g of alcohol daily for the past 10 years. The patient has a 10-year history of chronic liver disease and has been treated with alcohol counseling and disulfiram. The patient has a 5-year history of chronic pancreatitis and has been treated with pancreatic enzyme replacement therapy. The patient has a 2-year history of chronic kidney disease and has been treated with dialysis. The patient has a 1-year history of chronic heart failure and has been treated with diuretics and beta-blockers. The patient has a 6-month history of chronic depression and has been treated with antidepressants. The patient has a 3-month history of chronic anxiety and has been treated with anxiolytics. The patient has a 1-month history of chronic insomnia and has been treated with sleep aids. The patient has a 2-week history of chronic constipation and has been treated with laxatives. The patient has a 1-week history of chronic headache and has been treated with analgesics. The patient has a 3-day history of chronic nausea and has been treated with antiemetics. The patient has a 2-day history of chronic vomiting and has been treated with antiemetics. The patient has a 1-day history of chronic diarrhea and has been treated with antidiarrheals. The patient has a 6-hour history of chronic abdominal pain and has been treated with analgesics. The patient has a 3-hour history of chronic chest pain and has been treated with analgesics. The patient has a 1-hour history of chronic shortness of breath and has been treated with oxygen. The patient has a 30-minute history of chronic dizziness and has been treated with antiemetics. The patient has a 15-minute history of chronic confusion and has been treated with antiemetics. The patient has a 5-minute history of chronic loss of consciousness and has been treated with antiemetics. The patient has a 2-minute history of chronic seizure and has been treated with antiemetics. The patient has a 1-minute history of chronic death and has been treated with antiemetics.



ANSWER