



Code: 9207 NPN: FOOD

Format: Powder Size: 1 kg

PGX®

Weight Loss Meal Replacement French Vanilla · 1 kg

- A delicious and satisfying meal replacement that curbs appetite and prevents food cravings
- Low glycemic load and index; helps normalize postprandial blood glucose levels
- High in protein and suitable for carbohydrate-reduced diets
- · Convenient breakfast and/or lunch meal-replacement alternative
- Contains 5 g of PGX per serving
- No artificial sweeteners or flavours
- Also available in Double Chocolate

PRODUCT SUMMARY

PGX Weight Loss Meal Replacement promotes healthy weight management by reducing postprandial hyperglycemia, increasing satiety, and providing supportive nutrients for blood sugar metabolism. PGX (PolyGlycopleX®) is a highly viscous combination of well purified water-soluble polysaccharides derived from konjac root, sodium alginate, and xanthan gum, which has been shown to reduce postprandial glucose concentrations and lower total and LDL cholesterol, primarily by forming a gel matrix which slows gastric emptying. Postprandial hyperglycemia is an important causal factor in endothelial dysfunction and weight gain. By reducing the glycemic index of foods, viscous fibres prevent postprandial spikes in blood glucose concentration and in insulin production. They have also been found to increase satiety and reduce appetite, promoting a lower caloric intake, resulting in weight loss when combined with an energy-reduced diet.

PGX is combined with whey protein and medium chain triglycerides, shown clinically to enhance body fat loss while preserving lean muscle and to induce satiety and a decreased consumption of calories, respectively. A full complement of vitamins and minerals ensure support for healthy macronutrient metabolism.



To Place Your Order Email: customerservice@bioclinicnaturals.com **Call:** 1·888·826·9625 • **Fax:** 1·877·433·9862





WEIGHT LOSS MEAL REPLACEMENT · FRENCH VANILLA · 1 KG



Serving Size: 2 scoops (approx. 62 g)

Servings Per Container: 16

Key Ingredients:

Whey Protein Concentrate and Isolate (Microfiltered, Undenatured)	32 g
PGX® (PolyGlycopleX®)	5 g
Highly Purified Fibre Complex Manufactured Using the Proprietary EnviroSimplex® Process: Konjac-mannan (Amorphophallus konjac) (root), Sodium Alginate, Xanthan Gum	
Soy Lecithin	5.7 g
Medium Chain Triglycerides	

Ingredients: Whey protein concentrate and isolate (milk) (microfiltered, undenatured), Soy lecithin, PGX® polysaccharides complex (glucomannan, xanthan gum, sodium alginate), Organic cane sugar, Xylitol, Modified palm oil (medium chain triglycerides), Dicalcium phosphate dihydrate, Potassium citrate, Magnesium citrate, Sodium chloride, Calcium citrate, Ascorbic acid, Zinc citrate, Niacinamide, Biotin, Ferrous fumarate, *d*-alpha Tocopheryl acetate, Copper chelate, Beta-carotene, Molybdenum citrate, Selenium chelate, Manganese citrate, Calcium pantothenate, Folic acid, Thiamine hydrochloride, Pyridoxine hydrochloride, Riboflavin, Vitamin B12, Potassium iodide, Chromium chelate, Vitamin D2, Natural flavours.

Suggested Usage: To make a delicious and satisfying meal replacement smoothie, blend, shake, or briskly stir 2 scoops (approx. 62 g) of PGX® Weight Loss Meal Replacement into 350–500 mL of water 1–2 times per day or as directed by a health care practitioner. Drink additional water (350–500 mL) following the smoothie.

When first taking PGX® Weight Loss Meal Replacement, begin with 1 scoop (approx. 31 g) in a glass of water (175–350 mL) and gradually increase to 2 scoops (approx. 62 g) as directed above.

Caution: If you are under 18 years of age, are pregnant or breastfeeding, or have any health concerns, consult your health care practitioner before starting any weight loss plan. If you are taking medication, take it 1 hour prior to or 2 hours after taking PGX®. Keep out of reach of children.

Drug Interactions: A reduction in serum glucose concentrations may require a lowering of the dosage of any antidiabetic medications, and serum glucose should be monitored closely to prevent hypoglycemia. Oral medications may need to be taken one hour prior or two hours after to prevent a slowing down in their absorption. Similarly, fat-soluble vitamins should be taken separately. Theoretically, whey protein may decrease levodopa absorption and should be taken together with caution.

Contains no artificial colours, preservatives, or sweeteners; no wheat, gluten, yeast, egg, fish, shellfish, tree nuts, or GMOs. Suitable for vegetarians. Sealed for your protection. Do not use if seal is broken. For freshness, store in a cool, dry place.

References available at bioclinicnaturals.com



· GUARANTEED ·

Bioclinic Naturals* products are manufactured to meet or exceed current Good Manufacturing Practices (cGMP) of the U.S. Food and Drug Administration (FDA), Health Canada, and the Therapeutic Goods Administration (TGA) of Australia.





PRODUCT OF CANADA Bioclinic Naturals® Canada Burnaby, BC V3N 4T6 bioclinicnaturals.com