

QUESTION

1. A patient with a long history of alcohol abuse is brought to the emergency department with a 2-day history of confusion and vomiting.

On physical examination, the patient is found to have tachycardia, hyperreflexia, and rigidity of the neck.

Which of the following is the most likely diagnosis?

A. Acute alcohol withdrawal

B. Delirium due to alcohol intoxication

C. Delirium due to alcohol withdrawal

D. Delirium due to alcohol poisoning

E. Delirium due to alcohol withdrawal and alcohol poisoning

F. Delirium due to alcohol withdrawal and alcohol poisoning

G. Delirium due to alcohol withdrawal and alcohol poisoning

H. Delirium due to alcohol withdrawal and alcohol poisoning

I. Delirium due to alcohol withdrawal and alcohol poisoning

J. Delirium due to alcohol withdrawal and alcohol poisoning

K. Delirium due to alcohol withdrawal and alcohol poisoning

L. Delirium due to alcohol withdrawal and alcohol poisoning

M. Delirium due to alcohol withdrawal and alcohol poisoning

N. Delirium due to alcohol withdrawal and alcohol poisoning

O. Delirium due to alcohol withdrawal and alcohol poisoning

P. Delirium due to alcohol withdrawal and alcohol poisoning

Q. Delirium due to alcohol withdrawal and alcohol poisoning

R. Delirium due to alcohol withdrawal and alcohol poisoning

S. Delirium due to alcohol withdrawal and alcohol poisoning

T. Delirium due to alcohol withdrawal and alcohol poisoning

U. Delirium due to alcohol withdrawal and alcohol poisoning

V. Delirium due to alcohol withdrawal and alcohol poisoning

W. Delirium due to alcohol withdrawal and alcohol poisoning

X. Delirium due to alcohol withdrawal and alcohol poisoning

Y. Delirium due to alcohol withdrawal and alcohol poisoning

Z. Delirium due to alcohol withdrawal and alcohol poisoning

ANSWER

1. A patient with a long history of alcohol abuse is brought to the emergency department with a 2-day history of confusion and vomiting.

On physical examination, the patient is found to have tachycardia, hyperreflexia, and rigidity of the neck.

Which of the following is the most likely diagnosis?

A. Acute alcohol withdrawal

B. Delirium due to alcohol intoxication

C. Delirium due to alcohol withdrawal

D. Delirium due to alcohol poisoning

E. Delirium due to alcohol withdrawal and alcohol poisoning

F. Delirium due to alcohol withdrawal and alcohol poisoning

G. Delirium due to alcohol withdrawal and alcohol poisoning

H. Delirium due to alcohol withdrawal and alcohol poisoning

I. Delirium due to alcohol withdrawal and alcohol poisoning

J. Delirium due to alcohol withdrawal and alcohol poisoning

K. Delirium due to alcohol withdrawal and alcohol poisoning

L. Delirium due to alcohol withdrawal and alcohol poisoning

M. Delirium due to alcohol withdrawal and alcohol poisoning

N. Delirium due to alcohol withdrawal and alcohol poisoning

O. Delirium due to alcohol withdrawal and alcohol poisoning

P. Delirium due to alcohol withdrawal and alcohol poisoning

Q. Delirium due to alcohol withdrawal and alcohol poisoning

R. Delirium due to alcohol withdrawal and alcohol poisoning

S. Delirium due to alcohol withdrawal and alcohol poisoning

T. Delirium due to alcohol withdrawal and alcohol poisoning

U. Delirium due to alcohol withdrawal and alcohol poisoning

V. Delirium due to alcohol withdrawal and alcohol poisoning

W. Delirium due to alcohol withdrawal and alcohol poisoning

X. Delirium due to alcohol withdrawal and alcohol poisoning

Y. Delirium due to alcohol withdrawal and alcohol poisoning

Z. Delirium due to alcohol withdrawal and alcohol poisoning