## Citrus Disinfectant

ACTIVE INGREDIENT:

## KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID: IF IN EYES: Hold evelids open and flush with a steady, gentle stream of water for 15 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a Poison Control Center or doctor for further treatment advice

Follow the directions for use on the pesticide label when applying this product. Further sale or distribution of this product is strictly prohibited.

Customer Affixed Label

EPA Reg. No. 34810-35-1658



BACTERICIDAL: contact time: Pseudomonas aeruginosa, Salmonella rinsing or wiping is required.

MRSE. Immunodeficiency Virus Type 1 (Strain HTLV-IIIB), remain visibly wet for at least 5 minutes. Wipe or let air dry. DECONTAMINATING SURFACES/OBJECTS SOILED Influenza A Virus (Strain Hong Kong), Respiratory No rinsing or wiping is required except on food contact WITH BLOOD AND BODY FLUIDS WHICH MAY Syncytial Virus. 3-minute contact time: Human surfaces which require a potable water rinse after use. CONTAIN HIV-1. Coronavirus, Rotavirus, Feline Calicivirus (surrogate for Pre-cleaning is required for disinfection of Human PERSONAL PROTECTION: When handling objects soiled Norovirus), Hepatitis B Virus (HBV), Hepatitis C Virus Coronavirus. (HCV), Rhinovirus. 5-minute contact time: Adenovirus FUNGICIDAL DIRECTIONS: This product is effective Virus Type 2, Vaccinia Virus.

Trichophyton interdigitale.

TUBERCULOCIDAL: 5-minute contact time: Mycobacterium bovis at 20°C / 68°F on previously cleaned hard non-porous inanimate surfaces

SANITIZER: NONFOOD CONTACT 1-minute contact time: Klebsiella pneumoniae. FOOD CONTACT SANITIZER: 1-minute contact time:

PRECAUTIONARY STATEMENTS

Staphylococcus aureus, Escherichia coli.

HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco, or using the

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. TO CLEAN/DEODORIZE: Apply undiluted to hard non-porous surfaces by spraying, wiping with a cloth or towel until clean.

In the presence of 5% blood serum, is effective against: TO SANITIZE: Food contact: Preclean. Wet the surface. devices prior to sterilization or high-level disinfection. contact time: Let stand for 60 seconds on hard, non-porous surfaces. \*KILLS HIV ON PRE-CLEANED ENVIRONMENTAL pneumoniae. Wipe or let air dry. Rinse surface with potable water after SURFACES/OBJECTS PREVIOUSLY SOILED WITH Enterococcus faecalis Vancomycin Resistant (VRE), use. Non-food contact: Preclean. Wet the surface. Let BLOOD/BODY FLUIDS in all settings in which there is an Staphylococcus aureus - MRSA, Escherichia coli. 5-minute stand for 60 seconds on hard, non-porous surfaces. No expected likelihood of soiling of hard, non-porous

Listeria monocytogenes, Staphylococcus epidermidis - with visible soil prior to disinfection. Apply the product can be associated with the potential for transmission of undiluted to hard, non-porous surfaces with a sprayer, human immunodeficiency virus Type 1 (HIV-1) (associated VIRUCIDAL\*: 1-minute contact time: SARS-CoV-2 cloth, mop or brush, or by immersing equipment to be with AIDS virus). (cause of COVID-19) (Strain USA-WA 1/2020), Human disinfected. Wet all surfaces thoroughly. The surface must SPECIAL INSTRUCTIONS FOR CLEANING AND

Type 2, Herpes Simplex Virus Type 1, Herpes Simplex against Trichophyton interdigitale (athlete's foot fungus and CLEANING PROCEDURE: Blood and other body fluids a cause of ringworm) on inanimate surfaces in locker FUNGICIDAL: 5-minute contact time: Candida albicans, rooms, dressing rooms, exercise facilities, and shower and before application of the disinfectant. bath areas. Follow disinfection directions.

AREAS OF USE: On washable hard, non-porous surfaces of Hospitals, Kitchens, Bathrooms, Gyms, Health Care Facilities, Medical Offices, Dental Offices, Day Care to kill all bacteria, viruses\* and fungi listed on the label. Centers, Nurseries, Schools, and Nursing Homes.

TYPES OF SURFACES: On hard non-porous surfaces of floors. walls. fixtures, furniture, metal, stainless steel, chrome, CuVerro® antimicrobial copper, glazed porcelain and glazed ceramic tile, vinyl, plastic, baked enamel, glass, infant care equipment, kitchen surfaces, wrestling mats, exercise equipment, radiology equipment surfaces, computer surfaces, and keyboards.

DECONTAMINATION OF CRITICAL OR SEMI-CRITICAL DEVICES PRIOR TO TERMINAL STERILIZATION/HIGH LEVEL DISINFECTION: This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or, (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical

surfaces/objects with blood or body fluids and in which the enterica, Staphylococcus aureus, Escherichia coli 0157:H7, TO DISINFECT: Product is ready to use. Preclean areas surfaces/objects likely to be soiled with blood or body fluids

with blood or body fluids use disposable latex gloves. gowns, mask and eve coverings.

must be thoroughly cleaned from surfaces and objects CONTACT TIME: Do Not Dilute. Precleaned surfaces

must remain visibly wet for one minute using mop, cloth or sponge to mitigate HIV-1 virus. Use a 5-minute contact time DISPOSAL OF INFECTIOUS MATERIAL: Blood and other fluids should be autoclaved and disposed of according to Federal, State and Local regulations for infectious waste disposal

## STORAGE AND DISPOSAL:

Store airtight at room temperature. Nonrefillable container. Do not reuse or refill this empty container. Offer empty container for recycling. If recycling is not available, discard