

QUESTION

1. A patient with a long history of alcohol abuse is brought to the emergency department with a 2-day history of confusion and tremors.

2. The patient has a 10-year history of alcohol abuse and has been drinking approximately 100 g of alcohol daily.

3. The patient has a 2-day history of confusion and tremors.

4. The patient has a 2-day history of confusion and tremors.

5. The patient has a 2-day history of confusion and tremors.

6. The patient has a 2-day history of confusion and tremors.

7. The patient has a 2-day history of confusion and tremors.

8. The patient has a 2-day history of confusion and tremors.

9. The patient has a 2-day history of confusion and tremors.

10. The patient has a 2-day history of confusion and tremors.

11. The patient has a 2-day history of confusion and tremors.

12. The patient has a 2-day history of confusion and tremors.

13. The patient has a 2-day history of confusion and tremors.

14. The patient has a 2-day history of confusion and tremors.

15. The patient has a 2-day history of confusion and tremors.

16. The patient has a 2-day history of confusion and tremors.

17. The patient has a 2-day history of confusion and tremors.

18. The patient has a 2-day history of confusion and tremors.

19. The patient has a 2-day history of confusion and tremors.

20. The patient has a 2-day history of confusion and tremors.

21. The patient has a 2-day history of confusion and tremors.

22. The patient has a 2-day history of confusion and tremors.

23. The patient has a 2-day history of confusion and tremors.

24. The patient has a 2-day history of confusion and tremors.

25. The patient has a 2-day history of confusion and tremors.

26. The patient has a 2-day history of confusion and tremors.

27. The patient has a 2-day history of confusion and tremors.

28. The patient has a 2-day history of confusion and tremors.

29. The patient has a 2-day history of confusion and tremors.

30. The patient has a 2-day history of confusion and tremors.

31. The patient has a 2-day history of confusion and tremors.

32. The patient has a 2-day history of confusion and tremors.

33. The patient has a 2-day history of confusion and tremors.

34. The patient has a 2-day history of confusion and tremors.

35. The patient has a 2-day history of confusion and tremors.

