

QUESTION

1. A patient with a long history of alcohol abuse is brought to the emergency department with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day. The patient is brought back to the emergency department 2 weeks later with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day.

Question	Answer	Explanation
1. A patient with a long history of alcohol abuse is brought to the emergency department with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day. The patient is brought back to the emergency department 2 weeks later with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day.	Alcohol withdrawal	The patient's symptoms are consistent with alcohol withdrawal. The patient's blood alcohol concentration is 0.25 g/dL, which is a level that can cause withdrawal symptoms. The patient's symptoms improve with lorazepam, which is a benzodiazepine used to treat alcohol withdrawal.

2. A patient with a long history of alcohol abuse is brought to the emergency department with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day. The patient is brought back to the emergency department 2 weeks later with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day.

ANSWER



EXPLANATION

1. A patient with a long history of alcohol abuse is brought to the emergency department with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day. The patient is brought back to the emergency department 2 weeks later with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient's vital signs are stable, and there are no other significant findings on physical examination. The patient is diagnosed with alcohol withdrawal and is started on lorazepam. The patient's symptoms improve, and the patient is discharged on the following day.