PRODUCT SPECIFICATIONS





Code: 9351 Size: 180 Enteripure[™] Softgels Actual Size: 25.11 mm x 9.94 mm



OptiMega-3®

Pharmaceutical Grade Molecularly Distilled · Ultra Purified

- Optimal 2:1 ratio EPA to DHA
- Pharmaceutical-grade omega-3 fish oil blend, standardized to meet and exceed USP standards for quality and consistency
- Targeted delivery using a technology that embeds pure and natural ingredients into the softgel casing to prevent them from breaking down in the stomach, ensuring no fishy aftertaste
- Safe from lipid peroxides and environmental pollutants, including heavy metals, pesticides, dioxins, PCBs, and other harmful compounds
- Sourced from wild anchovies, sardines, and/or mackerel, one of the best natural sources of EPA and DHA
- Harvested through sustainable fishing practices

PRODUCT SUMMARY

Omega-3 fatty acids from fish oil support a number of cardiovascular health factors, including supporting artery health, maintaining triglyceride levels already within the normal range, supporting blood pressure already within the normal range, and maintaining platelet counts already within the normal range.*¹⁻⁴ These fatty acids comprehensively support the maintenance of overall cardiovascular health.*^{5,6}

EPA and DHA also support cognitive function through multiple mechanisms, as they are necessary for neuronal membranes and thus normal neuronal function.^{*7,8} EPA and DHA are essential to organizing the body's response to cellular damage and the clearance of cellular debris, providing the substrates for prostaglandins, resolvins and protectins.^{*9-11}

Omega-3 fatty acids help support cardiovascular health, maintain healthy immunity, and support healthy neurological function.*¹²⁻¹⁷ EPA/DHA helps maintain normal neurocognitive function and support a healthy mood.*¹⁸⁻²⁰ Meta-analyses of randomized trials found supplementation helped maintain healthy lipid levels, support healthy metabolism, and maintain HbA1c already within the normal range.*²¹⁻²³



To Place Your Order Email: customersupport@bioclinicnaturals.com **Call:** 1.877.433.9860 • **Fax:** 1.877.433.9862





Supplement Facts



Other ingredients: Enteripure[™] softgel (gelatin, glycerin, purified water, pectin), natural vitamin E.

Serving Size: 1 Softgel

Servings Per Container: 180

Other Ingredients: Enteripure™ softgel (gelatin, glycerin, purified water, pectin), natural vitamin E.

Suggested Usage: 1 softgel 1–3 times per day or as directed by a health care professional.

Caution: Consult your health care professional prior to use if you are pregnant, trying to become pregnant, breastfeeding, taking medication, have a medical condition, or anticipate surgery. Keep out of reach of children.

Contraindications: Individuals with an allergy to fish or seafood should use caution, though fish oil is rarely allergenic. Both benefit and risk have been documented for those at risk of or being treated for cardiac arrhythmias, with close supervision indicated.²⁴ Pregnant and nursing women are often advised to consume a minimum of 300 mg of DHA per day. Although DHA is recognized as essential to neurological development, no suggested daily intakes have been made for children.

Drug Interactions: Fish oil may potentially reduce the need or dosage for blood pressure medications in some individuals. and patients should be closely monitored. Because fish oil may have an antithrombotic effect, caution has traditionally been advised for those on anticlotting, antiplatelet or anticoagulant medications, or those at high risk of bleeding. However, a multinational randomized and controlled trial found that fish oil did not increase perioperative bleeding.^{25,26} In amounts greater than 3 g per day, hyperglycemia has been observed in diabetics and those with hypertriglyceridemia and close monitoring of patients on antidiabetic medication is recommended.²⁷

Contains no artificial colors, preservatives, or sweeteners; no dairy, sugar, wheat, gluten, yeast, soy, corn, egg, shellfish, salt, tree nuts, or GMOs. Sealed for your protection. Do not use if seal is broken. For freshness, store in a cool, dry place.

References

- Pischon, T., Hankinson, S.E., Hotamisligil, G.S., et al. (2003). *Circulation, 108*(2), 155-60. Swanson, D., Block, R., Mousa, S.A. (2012). *Adv Nutr,* 3(1), 1-7. Wang, C., Harris, W.S., Chung, M., et al. (2006). *Am J Clin Nutr,* 84(1), 5-17.

- 4 Saito, Y., Yokoyama, M., Origasa, H., et al. (2008). Atherosclerosis, 200(1), 135-40. 5. Delgado-Lista, J., Perez-Martinez, P., Lopez-Miranda, J., et al. (2012). Br J Nutr, 107(S2), S201-13.
- Chareonrungrueangchai, K., Wongkawinwoot, K., Anothaisintawee, T., et al. (2020). 6. Nutrient, 12(4), 1088.
- Lin, P.Y., Chiu, C.C., Huang, S.Y., et al. (2012). J Clin Psychiatry, 73(9), 1245-54.
- Dyall, S.C. (2015). Front Aging Neurosci, 7, 52. Yedgar, S., Krimsky, M., Cohen, Y., et al. (2007). Trends Pharmacol Sci, 28(9), 459-64. 8
- Schmitz, G., & Ecker, J. (2008). Prog Lipid Res, 47(2), 147-55. 10.
- Molfino, A., Amabile, M.I., Monti, M., et al. (2017). Oxid Med Cell Longev, 2017, 5987082. Löfvenborg, J.E., Andersson, T., Carlsson, P.O., et al. (2014). Nutr Diabetes, 4(10), e139. 12.
- Olsen, S.F., Østerdal, M.L., Salvig, J.D., et al. (2008). Am J Clin Nutr, 88(1), 167-75
- 14. Proudman, S.M., James, M.J., Spargo, L.D., et al. (2015). Ann Rheum Dis, 74(1), 89-95.

- Romano, C., Cucchiara, S., Barabino, A., et al. (2005). World J Gastroenterol, 11(45), 15. 7118-21.
- Di Giuseppe, D., Wallin, A., Bottai, M., et al. (2014). Ann Rheum Dis, 73(11), 1949-53. 17. Lorente-Cebrián, S., Costa, A.G., Navas-Carretero, S., et al. (2015). J Physiol Biochem, 71(2), 341-9.
- 18. Chang, J.P., Su, K.P., Mondelli, V., et al. (2018). Neuropsychopharmacology, 43(3), 534-45. Zhang, X.W., Hou, W.S., Li, M., et al. (2016). Aging Clin Exp Res, 28(1), 165-6. Bae, J.H., & Kim, G. (2018). Nutr Res, 50, 1-9. 19.
- 20.
- Chewcharat, A., Chewcharat, P., Rutirapong, A., et al. (2020). PLoS One, 15(2), e0228315. 21.
- O'Mahoney, L.L., Matu, J., Price, O.J., et al. (2018). *Cardiovasc Diabetol*, *17*(1), 98. Gao, H., Geng, T., Huang, T., et al. (2017). *Lipids Health Dis*, *16*(1), 131. Kromhout, D., & de Goede, J. (2014). *Am J Cardiol*, *99*(6A), 44C-46C. 22.
- 23.
- 24.
- Akintoye, E., Sethi, P., Harris, W.S., et al. (2018). Circ Cardiovasc Qual Outcomes, 11(11), 25. e004584
- 26. Gao, H., Geng, T., Huang, T., et al. (2017). Lipids Health Dis, 16(1), 131.



· GUARANTEED ·

Bioclinic Naturals® products are guaranteed to meet or exceed Good Manufacturing Practices (GMP) of the U.S. Food and Drug Administration (FDA), Health Canada, and the Therapeutic Goods Administration (TGA) of Australia.



PRODUCT OF CANADA Manufactured for and distributed by Bioclinic Naturals® U.S. 14224 167th Ave. SE, Monroe WA, USA 98272 bioclinicnaturals.com

FOR PROFESSIONAL USE ONLY. *These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease. © All Rights Reserved Bioclinic Naturals® 2021. March 2, 2021. 9225761

Head office Assured Natural Distribution Inc., 104 - 3686 Bonneville Place, Burnaby, BC, Canada V3N 4T6 | U.S. Distribution office 14224 167th Avenue SE, Monroe, WA, USA 98272 Customer service 1.877.433.9860 · Fax 1.877.433.9862 · Email customersupport@bioclinicnaturals.com