

QUESTION

1. A patient with a long history of alcohol abuse presents with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is brought to the emergency department and is found to have a Glasgow Coma Scale score of 3. The patient is intubated and brought to the intensive care unit. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL.

ANSWER

The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL.

QUESTION

QUESTION	ANSWER
1. A patient with a long history of alcohol abuse presents with a severe headache and vomiting. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is brought to the emergency department and is found to have a Glasgow Coma Scale score of 3. The patient is intubated and brought to the intensive care unit. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL.	The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL.

ANSWERS

1. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL. The patient is found to have a blood alcohol concentration of 0.25 g/dL.