BIOADVANCED®
SCIENCE-BASED SOLUTIONS

Bermudagrass
Control For Lawns

CONCENTRATE

KILLS THE ROOT,
NOT THE LAWN



FAST-ACTING FORMULA
FOR VISIBLE RESULTS

TREATS
4,000
SQ. FT.

KEEP OUT OF REACH OF CHILDREN
CAUTION See Back Label for Additional Precautionary Statements

Active Ingredients:	% by wt.
Topramezone	0.104%
Sodium Salt of Bentazon	4.130%
Other Ingredients	95.766%
TOTAL	100.000%

THIS PRODUCT CONTAINS:
0.0088 lb. topramezone (CAS # 210631-68-8) per gallon or 0.104%
0.3205 lb. bentazon acid equivalent (acid of CAS # 50723-80-3) per gallon or 3.780%

5010000500 241120V1
NET CONTENTS 32 FL OZ (946ml)

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call 24/7 toll-free 1-877-229-3763 for medical emergency information.**

Active Ingredients: 0.104% Topramezone (CAS # 210631-68-8) and 4.130% Sodium Salt of Bentazon (CAS # 50723-80-3).

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic and terrestrial plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- These chemicals, topramezone and bentazon, have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- POLLINATOR ADVISORY STATEMENT: This product may adversely impact the forage and habitat of local pollinators, such as the monarch butterfly (and its larvae), birds, or bats. Protect wildlife by following label directions, and making only directed applications.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

BioAdvanced® is a registered trademark of SBM Life Science Corp.

PRESS TO RESEAL

5012000900 241120V1

EPA Reg. No. 92564-85 EPA Est. No indicated by 3rd and 4th digits of the batch number on this package. (75) = 5905-AR-1 (39) = 58996-MO-1 (57) = 071106-GA-003 (46) = 88746-MO-1 (42) = 88746-GA-01 (65) = 92564-TX-1 Produced For: SBM Life Science Corp. PO Box 525, Morrisville, NC 27560 BioAdvanced® is a registered trademark of SBM Life Science Corp.

BIOADVANCED
SCIENCE-BASED SOLUTIONS

Bermudagrass Control For Lawns

CONCENTRATE

KILLS/CONTROLS

BERMUDAGRASS

ALSO KILLS

CRABGRASS | FOXTAIL | CLOVER

JAPANESE STILTGRASS | PLUS OTHER WEEDS AS LISTED

See inside booklet for a complete list of weeds controlled

FOR USE ON

KENTUCKY BLUEGRASS | FESCUE | RYEGRASS

FOR QUESTIONS OR COMMENTS,
Call toll-free
1-877-229-3724 or visit
www.bioadvanced.com

SATISFACTION 100% GUARANTEED‡‡

‡‡*Or your money back.*

FOR OUTDOOR RESIDENTIAL USE ONLY

8 40216 20254 2

5012000900 241120V1

READ THE ENTIRE LABEL FIRST

Indicaciones en español después de las de inglés

Produced For: SBM Life Science Corp.
PO Box 525, Morrisville, NC 27560
BioAdvanced® is a registered trademark of SBM Life Science Corp.

PEEL HERE

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WHERE TO USE

ON ESTABLISHED LAWNS without harming lawn grasses – Kentucky Bluegrass, Ryegrass, Fescue

DO NOT apply to bahiagrass, buffalograss, carpetgrass, centipedegrass, clover (desirable), creeping bentgrass, dichondra, St. Augustinegrass (including Floratam), seashore paspalum or zoysiagrass.

HOW MUCH TO USE

Use as a broadcast treatment (treating whole areas of the lawn). Measure area to be sprayed. Calculate 'square feet' by multiplying length times width.

This bottle will treat 4,000 sq. ft. in about 20 minutes (actual time will vary depending on water pressure)

DO NOT apply more than 8 fl oz per 1,000 sq ft (equivalent to 0.02 lb topramezone and 0.87 lb bentazon acid per acre) per application.

DO NOT make more than two applications per year.

DO NOT apply more than 16 fl oz per 1,000 sq ft (equivalent to 0.04 lb topramezone and 1.75 lb bentazon acid per acre) per year.

HOW TO USE

Shake bottle well before use.

For first time use, remove plastic shipping lock.



CONNECT
1. Shake well before use.
2. Connect sprayer to garden hose.
3. Make sure that the Sprayer is in the OFF position.
4. Turn on water at faucet.

SPRAY
1. Extend hose to farthest area to be treated, point Sprayer nozzle in direction to be sprayed.
2. Turn Sprayer to ON position.
3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap treated areas. Work back toward faucet to avoid walking over wet, treated areas.

FINISH
1. To stop spraying, turn Sprayer to OFF position.
2. Turn off water at faucet.
3. Relieve water pressure in hose by turning Sprayer to ON position until water slows to a drip, then turn Sprayer to OFF position.
4. Disconnect Sprayer from hose.

FOR BEST RESULTS

WHEN TO APPLY

i. Spray when
   1. weeds are small and actively growing
   2. temperature is below 90°F

ii. To control Bermudagrass, apply in spring to late summer when Bermudagrass begins to green up. Up to two applications spaced three weeks apart may be needed for well-established Bermudagrass (do not make more than two applications per year).

iii. Certain hard-to-kill weeds may be retreated in 3 to 4 weeks if necessary.

iv. Bare spots may be seeded 4 weeks after application.

v. Treated fescue and perennial ryegrass lawns may be re-seeded immediately after application. However, wait 3 weeks after application for Kentucky bluegrass before reseeding.

vi. Before treating new or renovated lawns:
   - Apply only to well established seedlings or sprigs. New grass seedlings must be at least 1 month old (3 months for Kentucky bluegrass).
   - Kentucky bluegrass sod must be at least 1 month old since installation.

MOWING

Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING

If lawn is dry, water 1 to 2 days before using this product.

RESTRICTIONS

DO NOT:

- Spray or allow to drift on plants growing in or around the lawn such as trees, shrubs, flowers, perennials, groundcovers, vegetables, & herbs.
- Water or irrigate for 1 hour after application.
- Use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Apply while grass is emerging from dormancy.
- Mow within **2 days before or 2 days** after spraying.
- Enter or allow people or pets to enter the treated area until spray has dried. People and pets may enter treated areas after spray has dried.
- Apply this product in a way that will contact people either directly or through drift, or if winds speeds are greater than 10 mph.

(cont'd)

RESTRICTIONS *(cont'd)*

DO NOT:

- Apply more than 2 times per year.
- Apply for at least 28 days after seeding of tolerant turfgrasses. Seed prior to application and apply only to established seedlings and sprigs.
- Apply to drought stressed lawns and/or drought stressed weeds, exposed feeder roots of trees or ornamentals, or within the dripline of trees and other ornamental plants, or within ornamental beds, or to areas where runoff into those areas is likely to occur.
- Use around these ornamental plants: Bar Harbor juniper, philodendron, pittosporum, Podocarpus, salvia.
- Use clippings as mulch around flowers, ornamentals, trees or in vegetable gardens.
- Apply an organophosphate (such as malathion) or carbamate insecticide (such as carbaryl) within 1 week of applying this product or lawn injury may result.

HOW IT WORKS

This Product contains **Two** proven weed killers to control listed lawn weeds & crabgrass down to the root(s) without harming your lawn when used as directed.

WEEDS KILLED/CONTROLLED

Amaranth, Palmer
Amaranth, Powell
Anoda, Spurred
Balloon vine
Barnyard grass
Beggarticks
Bermudagrass, common
Bindweed (Field, Hedge)
Buckwheat, Wild
Burcucumber
Canada Thistle
Carpetweed
Chickweed, common
Clover, large hop
Clover, white
Cocklebur
Cocklebur, common
Crabgrass, large
Crabgrass, smooth
Crabgrass, southern
Croton, Tropic
Cupgrass, woolly
Dallisgrass
Dandelion, common
Dayflower
Devil's claw
Eclipta
Foxtail, giant
Foxtail, green
Foxtail, yellow
Goosegrass
Galinsoga
Galinsoga, hairy
Ground ivy
Groundsel, Common
Horseweed (Marestail)
Jimsonweed
Johnsongrass seedling
Kochia
Ladysthumb
Lambsquarters, common
Lawn Burweed
Mallow, common
Mallow, Venice
Marsh elder
Maywood/dogfennel
Millet, wild proso
Morning glory
Morning glory spp
Mustard spp
Mustard, Wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Nimblewill
Nutsedge, Yellow
Oxalis (Yellow wood sorrel)
Panicum, fall
Paspalum, slender
Pigweed, prostrate
Pigweed, redroot
Pigweed, smooth

(cont'd)

WEEDS KILLED/CONTROLLED *(cont'd)*

Pigweed, tumble
Poinsettia, Wild
Prickly lettuce
Purslane, Common
Radish, Volunteer
Ragweed, Common
Ragweed, giant
Redweed
Sedge, Annual
Senna, Coffee
Sesbania
Shattercane
Shepherd's-purse
Sida, Prick or Teaweed
Sida, prickly
Signalgrass, broadleaf
Smartweed
Smartweed, Pennsylvania
Speedwell (Veronica spp.)
Spurweed
Starbur, Bristly
Stiltgrass, Japanese
Sugar Beet, Volunteer
Sunflower, wild (common)
Thistle, Russian
Velvetgrass, common
Velvetleaf
Venice Mallow
Waterhemp
Windmill grass
Zoysiagrass



Do not apply this product in a way that will contact any person or pet either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until spray has dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING

If Empty:

- Non-refillable container. Do not refill or reuse this container.
- Place in trash or offer for recycling if available.

If Partly Filled:

- Call your local solid waste agency for disposal instructions.
- Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call 24/7 toll-free 1-877-229-3763 for medical emergency information.**

Active Ingredients: 0.104% Topramezone (CAS # 210631-68-8) and 4.130% Sodium Salt of Bentazon (CAS # 50723-80-3).

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

- This product is toxic to aquatic and terrestrial plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- These chemicals, topramezone and bentazon, have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- POLLINATOR ADVISORY STATEMENT: This product may adversely impact the forage and habitat of local pollinators, such as the monarch butterfly (and its larvae), birds, or bats. Protect wildlife by following label directions, and making only directed applications.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

INSTRUCCIONES DE USO

Constituye una violación a la ley federal usar este producto de manera diferente a lo establecido en esta etiqueta.

DÓNDE USAR

EN CÉSPEDES YA ESTABLECIDOS sin dañar el pasto – césped azul de Kentucky, raigrás, festuca

No aplicar al césped bahía, césped búfalo, pasto chato, césped ciempiés, trébol (si lo desea mantener), Agrostis stolonifera, dichondra, césped San Agustín (incluyendo Floratam), césped paspalum de la costa y césped zoysia

CUÁNTO USAR

Usar como tratamiento por esparcimiento (tratamiento de todas las áreas del césped). Medir el área a rociar. Calcular los pies cuadrados multiplicando la longitud por el ancho.

Esta botella sirve para tratar hasta 4,000 pies cuadrados en alrededor de 20 minutos (el tiempo real puede variar dependiendo de la presión de agua)

NO aplicar más de 8 onzas líqui. (Equivalente a 0.02 libras de topramezone y 0.87 libras de ácido de bentazón por acre) por aplicación.

NO realizar más de dos aplicaciones por año.

NO aplicar más de 16 onzas líqui.por 1,000 pies cuadrados (equivalente a 0.04 libras de topramezone y 1.75 libras de ácido de bentazón por acre) por año.

CÓMO USAR

Agitar bien el concentrado antes de usar.

Cuando se use por primera vez, quitar el bloqueo de plástico de envío.



CONECTAR
1. Agitar bien antes de usar.
2. Conectar el rociador de manera segura a la manguera del jardín.
3. Asegúrese que el Rociador esté en posición cerrado OFF.
4. Abrir el grifo de agua.

ROCIAR
1. Extender la manguera hasta el área más lejana, apuntar la boquilla del rociador en dirección hacia donde desea rociar.
2. Girar el rociador a la posición ON.
3. Caminar a un ritmo constante mientras rocía, usando un movimiento oscilante y volver a regar la misma área tratada solo un poco. Caminar hacia atrás hacia el grifo para evitar caminar sobre el césped húmedo y tratado.

TERMINAR
1. Para finalizar el rocío, llevar el rociador hasta la posición OFF.
2. Apagar el grifo de agua.
3. Liberar la presión de agua en la manguera girando la válvula del rociador nuevamente a la posición ON hasta que el agua disminuya lentamente a un goteo, luego volver a girar la válvula del rociador a la posición OFF.

PARA MEJORES RESULTADOS

CUÁNDO APLICAR

i. Rociar cuando
   1. Las malezas son pequeñas y están creciendo activamente
   2. La temperatura sea debajo de los 90°F

ii. Para controlar el césped Bermuda, aplicar entre la primavera hasta fines del verano cuando el césped Bermuda comienza a ponerse verde. Puede ser necesario hacer hasta dos aplicaciones espaciadas entre tres semanas para el césped Bermuda bien establecido (no hacer más de dos aplicaciones por año).

iii. Ciertas malezas que son difíciles de matar pueden volver a tratarse en 3 a 4 semanas si fuese necesario.

iv. Las manchas sin nada pueden sembrarse en 4 semanas luego de su aplicación.

v. Los céspedes raigrás perenne y festuca que hayan sido tratados pueden volver a sembrarse inmediatamente después de la aplicación. Pero, espere 3 semanas después de la aplicación antes de plantar el césped azul de Kentucky.

vi. Antes de aplicar a césped nuevo o renovado:
   • Aplicar solamente a gajos o plántulas bien establecidos. Las nuevas plántulas de césped deben tener por lo menos 1 mes (3 meses para el césped azul Kentucky)
   • El césped azul de Kentucky debe tener como mínimo 1 mes desde su instalación.

CORTADO

Los recortes de los tres primeros cortes deben dejarse en las áreas tratadas. No usar los restos de cortes de las áreas tratadas como mantillo o compost alrededor de flores, plantas decorativas, árboles o en jardines de vegetales.

REGADO

Si el césped está seco, riegue 1 a 2 días antes de usar este producto.

RESTRICCIONES

NO:

- Rociar o permitir que gotee en plantas que crecen en o alrededor de plantas tales como arboles, arbustos, flores, plantas perennes, plantas de cobertura, verdura o hierbas.
- Regar o irrigar durante 1 hora después de la aplicación.
- Usar en césped que se usa para la venta u otro uso comercial como pasto, o para producción comercial de semillas o con fines de investigación.
- Aplicar cuando el césped está emergiendo del letargo.
- Cortar **2 días antes o 2 días** después de rociar.
- Ingresar o permitir que las personas y las mascotas vuelvan a ingresar al área tratada hasta que el rocío se haya secado. Las personas y los animales pueden volver a ingresar al área tratada luego de que el rocío se haya secado.
- Aplicar este producto en un modo que entre en contacto con las personas directamente o por goteo, o si la velocidad del viento es mayor a 10 mph.

(cont'd)

RESTRICCIONES *(cont'd)*

NO:

- Aplicar más de 2 veces al año.
- Aplicar durante por lo menos 28 días después de sembrar hierba de césped que lo tolere. Sembrar antes de aplicar y aplicar solo en gajos o plántulas bien establecidos.
- Aplicar a céspedes estresados por sequía o malezas estresadas por sequía, raíces expuestas de árboles o plantas decorativas o dentro de la línea de riego de árboles o plantas decorativas o en áreas donde puede correr el agua hacia esa área.
- Usar alrededor de plantas decorativas: enebro rastrero junípero, philodendron, pitosporo, pino de cerro, salvia.
- Usar los restos de cortes como el mantillo alrededor de flores, plantas ornamentales, árboles o huertas de verduras.
- Aplicar un organofosfato (tal como malatión) o insecticida carbamato (tal como carbaryl) dentro de 1 semana de aplicar este producto o puede causar daño al césped.

CÓMO FUNCIONA

Este producto contiene **dos** exterminadores probados de malezas para controlar el zácate cangrejo y las malezas del césped que aparecen en la lista hasta la raíz sin dañar el césped cuando se usa como se indica.

MALEZAS CONTROLADAS/EXTERMINADAS

Amaranth, Palmer
Amaranto, palmera
Amaranto, Powelli
Anoda cristata
Cardiospermum halicacabum
Césped Barnyard
Bidens frondosa
Césped Bermuda, común
Corregúela menor de campo
Buckwheat, salvaje
Sicyos angulatus
Cardo de Canadá
Axonopus
Oreja de ratón, común
Trébol amarillo grande
Trébol, blanco
Bardana menor
Bardana menor común
Zácate cangrejo, grande
Zácate cangrejo, de verano
Zácate cangrejo, del sur
Crotón, trópico
Pasto lanoso
Heno lenoso
Diente de león común
Commelina communis
Garra del diablo
Eclipta
Cola de zorro, gigante
Cola de zorro, verde
Cola de zorro, amarilla
Goosegrass
Galinsoga
Galinsoga, lanosa
Hiedra rastrera
Senecio común
Hierba carnicera (Marestail)
Higuera loca
Sorgo de alepo (plántula)
Kochia
Persicaria
Lambsquarters, común
Burweed engomada del césped
Malva, común
Malva, Venecia
Marsh elder
Mala madre/Hinojo de perro
Hierba de mijo silvestre
Gloria de la mañana salvaje
Gloria de la mañana, spp
Mostaza, spp
Mostaza, salvaje
Solanácea, negra
Solanácea, negra del este negro
Solanácea, peluda
Nimblewill
Juncia real, amarilla

(cont'd)

MALEZAS CONTROLADAS/EXTERMINADAS *(cont'd)*

Oxalis (madera amarilla)
Panicum, otoño
Paspalum, slender
Amaranto, próstata
Amaranto, raíz roja
Amaranto, suave
Amaranto, tumble
Poinsettia, salvaje
Lechuga espinosa
Verdolaga común
Rabanito, voluntario
Ambrosía común
Ambrosía , gigante
Secuoya roja
Junco, anual
Senna, café
Sesbania
Sorgo
Bolsa del pastor
Sida, espinosa o teaweed
Sida, espinosa
Césped señal de hoja ancha
Persicaria
Persicaria, Pennsylvania
Speedwell (Verónica spp.)
Espuela de gallo
Torito, Bristly
Hiedra zancuda, japonesa
Remolacha azucarera, voluntaria
Girasol, salvaje (común)
Cardo ruso
Velvetgrass, común
Malva terciopelo
Malva de Venecia
Amaranto de frutos ásperos
Chloris
Césped zoysia



No aplicar este producto de modo que entre en contacto con personas o mascotas ya sea directamente o por goteo. Mantener a las personas y mascotas fuera del área durante su aplicación. No permitir que las personas y las mascotas vuelvan a ingresar al área tratada hasta que el rocío se haya secado.

ALMACENAMIENTO Y ELIMINACIÓN

No contaminar alimentos, agua o alimentos para mascotas por almacenamiento o eliminación.

ALMACENAMIENTO DEL PESTICIDA

Almacenar en el contenedor original en un lugar seco y fresco, fuera del alcance de los niños, preferentemente en un gabinete de almacenamiento cerrado. Proteger de congelamiento.

ELIMINACIÓN DEL PESTICIDA Y MANIPULACIÓN DEL CONTENEDOR

Si está vacío:

- Contenedor no rellenable. No rellenar o volver a usar este contenedor.
- Colocar en la basura u ofrecer para su reciclado si fuese posible.

Si está parcialmente lleno:

- Llamar a la agencia de eliminación de desechos sólidos para conocer las indicaciones de eliminación.
- Nunca colocar el producto sin usar en un drenaje de salida o de entrada.

AVISO: Las investigaciones y pruebas han determinado que las “Indicaciones de uso” corresponden al uso adecuado de este producto en condiciones esperadas. Hasta el punto en que aplica la ley, el comprador asume la responsabilidad de falta de rendimiento o seguridad si no se usa de acuerdo con las instrucciones.

Garantizamos la devolución del dinero: Si no está satisfecho con este producto, le devolveremos el dinero de la compra original.

PRIMEROS AUXILIOS

SI LO TRAGA

- Llamar al centro de control de envenenamiento o al médico inmediatamente para asesoramiento sobre el tratamiento.
- Hacer que la persona sorba un vaso de agua si puede tragar.
- No inducir el vómito a menos que se lo indique un médico o el centro de control de envenenamiento.
- No dar nada por boca a una persona inconsciente.

NOTA: Cuando llame al centro de control de envenenamiento, lleve la etiqueta de este producto. Si acude en busca de tratamiento a una sala de emergencia o al consultorio de un médico, lleve esta etiqueta del producto para mostrarle al personal médico. **Puede llamar 24/7 a la línea gratuita 1-877-229-3763 para información sobre atención médica de urgencia.**
Ingredientes activos: 0.104% Topramezone (CAS # 210631-68-8) y 4.130% sal de sodio de Bentazón (CAS # 50723-80-3).

DECLARACIONES DE PRECAUCIÓN

Peligroso para animales domésticos y humanos

PRECAUCIÓN

- Peligroso si lo traga.
- Lavar cuidadosamente con agua y jabón luego de manipularlo y antes de comer, beber, masticar goma de mascar, usar tabaco o ir al baño.

PELIGROS AMBIENTALES

- Este producto es tóxico para las plantas acuáticas y de tierra.
- Para proteger al medio ambiente, no permitir que el pesticida ingrese o corra por los drenajes de tormenta, drenaje de aguas pluviales, canaletas o aguas superficiales. Aplicar este producto en clima calmo cuando no haya pronóstico de lluvia en las próximas 24 horas para asegurar que el viento o la lluvia no soplen o laven el pesticida fuera del área de tratamiento. Enjuagar el equipo usado para la aplicación en el área tratada ayudará a evitar que el producto ingrese en cuerpos de agua o sistemas de drenaje.
- Estos químicos, topramezone y bentazón, tienen propiedades y características asociadas a químicos detectados en las aguas subterráneas. Estos químicos pueden escurrirse a las aguas subterráneas si se usan en suelos donde sea permeable especialmente donde la mesa de agua sea poco profunda.
- AVISO DE AGENTES POLINADORES: Este producto puede tener un impacto adverso en el forraje y el hábitat local de los polinizadores locales, tales como la mariposa monarca (y su larva), los pájaros o los murciélagos. Es importante proteger la vida silvestre si sigue las instrucciones de la etiqueta, y hace solo las aplicaciones indicadas.

PELIGROS QUÍMICOS O FÍSICOS

No mezclar o tener contacto con el agente oxidante. Puede ocurrir una reacción química peligrosa.