
BIOADVANCED®
SCIENCE-BASED SOLUTIONS

SEASON LONG WEED KILLER

CONCENTRATE



KILLS LAWN WEEDS

PREVENTS WEEDS*
+ CRABGRASS UP TO 4 MONTHS

USE ON ALL COMMON LAWNS

FOR RESIDENTIAL OUTDOOR USE ONLY
KEEP OUT OF REACH OF CHILDREN
CAUTION See back booklet for additional precautionary statements
*Listed

Active Ingredients:	% by wt.
Dithiopyr	1.10%
Dicamba	0.44%
Penoxsulam	0.10%
Other Ingredients	98.36%
Total	100.00%

THIS PRODUCT CONTAINS:
0.092 lb. dithiopyr (CAS # 97886-45-8) per gallon
0.037 lb. dicamba (CAS # 1918-00-9) per gallon
0.0082 lb. penoxsulam (CAS# 219714-96-2) per gallon

NET CONTENTS
32 FL OZ (946 mL)
XXXXXXXX XXXXXXX

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.

NOTE: When calling a poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call 24/7 toll-free 1-877-229-3763 for medical emergency information.**

Active Ingredients: 0.44% Dicamba (CAS # 1918-00-9), 0.10% Penoxsulam (CAS# 219714-96-2), and 1.10% Dithiopyr (CAS # 97886-45-8)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

• Harmful if swallowed. • Causes moderate eye irritation. • Avoid contact with eyes, skin, or clothing. • After using this product wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

• This product is toxic to fish and highly toxic to aquatic invertebrates including oysters and shrimp. • To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. • Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. • Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Ground Water Advisory: This product has properties and characteristics associated with chemicals detected in groundwater. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. May result in ground water contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Non-Target Organism Advisory: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

EPA Reg. No. 92564-88 EPA Est. No indicated by 3rd and 4th digits of the batch number on this package. (75) = 5905-AR-1 (39) = 58996-MO-1 (57) = 071106-GA-003
Distributed by: SBM Life Science Corp. PO Box 525, Morrisville, NC 27560 BioAdvanced® is a registered trademark of SBM Life Science Corp.

PRESS TO RESEAL

XXXXXXX XXXXXXX

BIOADVANCED®
SCIENCE-BASED SOLUTIONS

SEASON LONG WEED KILLER

CONCENTRATE

KILLS

DANDELION | NUTSEDGE | POA ANNUA

CHICKWEED[1] | CLOVER | CARPETWEED[1]

PREVENTS

DANDELION | CRABGRASS | POA ANNUA

CHICKWEED[1] | CLOVER | CARPETWEED[1]

WHERE TO USE - ON ESTABLISHED LAWNS

COOL-SEASON GRASSES
Bentgrass, Kentucky Bluegrass, Fescue (Chewing, Creeping Red, Tall), Perennial Ryegrass

WARM-SEASON GRASSES
Bermudagrass (common or hybrid), Tall Fescue, Kikuyugrass, Zoysiagrass, Centipedegrass, St. Augustinegrass, Buffalograss

DO NOT USE ON ANY OTHER TURF TYPES INCLUDING:
dichondra, carpetgrass, bahiagrass, or turfgrass containing desirable clovers.

[1]Will be controlled or suppressed by this product when applied before weeds germinate

STOP! READ THIS ENTIRE LABEL FIRST

Indicaciones en español después de las de inglés

FPO

8 40216 20085 2

XXXXXXX XXXXXXX

FOR QUESTIONS OR COMMENTS,
Call toll-free 1-877-229-3724 or visit us online at www.bioadvanced.com

SATISFACTION 100% GUARANTEED#

#Or your money back.

FPO

Use your phone to scan this code to learn

PEEL HERE

QUICK FACTS

TWO-WAY ACTION

Kills & Prevents listed Weeds.

KILLS ON CONTACT

Use with confidence on most turf types including Kentucky bluegrass, perennial bluegrass, fescue, perennial ryegrass, bermudagrass, centipedegrass and St. Augustine grass (including floratam).
Takes the guesswork out of weed control in the lawn. For a beautiful, weed-free lawn.

COVERAGE

Covers 4000 sq. ft. of lawn
For questions or comments, call toll-free 1-877-229-3724 or visit our website at www.BioAdvanced.com

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE USE

Read and follow all label directions.

RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
- Keep people and pets out of the area during application and until spray has dried.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not apply directly to, or allow spray drift on flowers, ornamental shrubs, trees, or other desirable plants, as injury may occur.
- Do not apply within 10 feet from sensitive plants including vegetable gardens.
- Clippings from mowings should be left on the treated area. Do not use the clippings from the treated area as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not treat lawns under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury due to insects, nematodes or disease.
- Do not allow the product to enter any drain during or after application.
- Do not apply when rainfall is expected within 24 hours (for control minimum 1 hour before the product has sufficient time to dry).
- Do not make applications during rain.
- Do not apply under windy conditions that would cause the product to drift to sensitive plants.

(cont'd)

- Do not spray the product into fish pools, ponds, streams, or lakes.
- Do not apply more than 1.5 lb ai/A of dithiopyr per year (when using additional products containing dithiopyr).
- Do not apply more than 8 fl oz of this product per 1,000 sq. ft. per application (equivalent to 0.25 lb dithiopyr, 0.10 lb dicamba, and 0.02 lb penoxsulam per acre).
- Do not apply more than 4 times per year
- Do not exceed 32 fl oz product per 1,000 sq. ft. per year (equivalent to 1 lb dithiopyr, 0.40 lb dicamba, and 0.08 lb penoxsulam per acre).
- In the state of New York do not exceed 8 fl oz of product per 1,000 sq. ft. per year (equivalent to 0.25 lb of dithiopyr per acre per year)

PRECAUTIONS

- Use this product only on established lawns.
- Excessive rainfall within 24 hours after application may cause unintended run-off of pesticide application.
- Apply when the air is calm to avoid spray drift onto desirable plants.

WHEN TO USE

Apply anytime from spring through fall to actively growing established turf when temperatures are between 50-90 degrees F to kill broadleaf weeds. This product will not burn your lawn when temperatures rise, up to 90 degrees F.

WHERE TO USE

ON ESTABLISHED LAWNS

Cool Season Grasses: bentgrass, Kentucky bluegrass, fescue (chewing, creeping red, tall) and perennial ryegrass

Warm Season Grasses: bermudagrass (common or hybrid), tall fescue, kikuyugrass and zoysiagrass, centipedegrass, St. Augustinegrass, buffalograss.

DO NOT USE ON ANY OTHER TURF TYPES including: dichondra, carpetgrass, bahiagrass, or turfgrass containing desirable clovers.

HOW MUCH TO USE

Use as a broadcast treatment (treating whole areas of the lawn).This bottle treats up to 4,000 sq. ft. Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. The maximum application rate of this product is 8 fl. oz. per 1,000 sq. ft. per application. Do not apply more than 4 times per year (except in the state of New York). See Restrictions above for more information on maximum use rates.

HOW TO USE

To kill weeds, thorough lawn and weed coverage is necessary.

HOW TO ATTACH AND SPRAY

For first time use, remove plastic shipping lock.



CONNECT
1. Shake well before use.
2. Connect sprayer to garden hose.
3. Make sure that the Sprayer is in the OFF position.
4. Turn on water at faucet.

SPRAY
1. Extend hose to farthest area to be treated, point Sprayer nozzle in direction to be sprayed.
2. Turn Sprayer to ON position.
3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap treated areas. Work back toward faucet to avoid walking over wet, treated areas.

FINISH
1. To stop spraying, turn Sprayer to OFF position.
2. Turn off water at faucet.
3. Relieve water pressure in hose by turning Sprayer to ON position until water slows to a drip, then turn Sprayer to OFF position.
4. Disconnect Sprayer from hose.

HOW OFTEN

This product may be reapplied no sooner than 35 days after the previous application.

HOW IT WORKS

This product contains three selective herbicides – it selectively kills weeds in the lawn but won't harm the lawn itself.

FOR BEST RESULTS

WHEN TO APPLY

Spray when weeds are present. • Temperatures are between 50 - 90° F
• Do not apply while grass is emerging from dormancy.

MOWING

Do not mow for 1 to 2 days after spraying. • Clippings from mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING

If lawn is dry, water 1 to 2 days before using this product.

HOW THIS PRODUCT WORKS

This product contains a proven blend of two post emergent weed killers to kill existing weeds, plus a pre-emergent to prevent new weeds. And, it won't harm your lawn when used as directed.

Dicamba – Dicamba is a selective herbicide used to control a wide spectrum of broadleaf weeds. It has a broad application window and assists in the burn-down of weeds. Dicamba is rapidly absorbed by plant leaves, stems, and roots, and works quickly to disrupt weed growth. The maximum single application rate is equivalent to 0.10 lb ai/a.
Penoxsulam – Penoxsulam is a systemic herbicide that is absorbed by the leaves and roots of plants. Weeds treated with penoxsulam will stop growing almost immediately and die within 2 to 4 weeks after application. The maximum single application rate is equivalent to 0.02 lb ai/a.
Dithiopyr – Dithiopyr is a pre-emergent herbicide that keeps grassy and broadleaf weeds from germinating. It has post-emergent control of seedling crabgrass for up to 4 weeks after the crabgrass germinates and emerges. If crabgrass begins to spread (tiller), this product will not be effective and should not be applied later than 4 weeks after germination. The maximum single application rate is equivalent to 0.25 lb ai/a.
Like all weed killers, the speed of action is dependent on many conditions (e.g. weed type, temperature) at any given time.

KILLS BROADLEAF WEEDS

Alkanet
Amaranth, palmer
Amaranth, powell
Amaranth, spiny
Aster, slender
Aster, spiny
Aster, whiteheath
Bedstraw
Beggarweed, creeping
Beggarweed, dwarf
Betony, common
Bindweed, field
Bindweed, hedge[1]
Black medic[1]
Broomweed, common
Buckwheat, tartary, wild
Burclover, California
Burdock, common
Burnweed, America
Buttercup, corn
Buttercup, creeping
Buttercup, hairy
Buttercup, red-horn
Buttercup, rough weed
Buttercup, western field
Carpetweed[1]
Carrot, wild
Catchfly
Catchweed
Celery, wild
Chamomile, corn
Chervil, bur
Chickweed (common)[1]
Chickweed (mouseear)[1]
Clover
Clover, Dutch
Clover, honey suckle
Clover, hop
Clover, red
Clover, white
Cockle, corn, cow, white
Cocklebur, common
Coffeeweed (common chickory)
Copperleaf, hophornbeam
Cornflower (bachelor button)
Croton, tropic
Croton, wooly
Cudweed, purple
Curly dock
Cutleaf
Daisy, English
Dandelion, common
Dandelion, false
Dollarweed (pennywort)
Dragonhead, American
Eclipta
Evening primrose, cutleaf
Falseflax, smallseed
Fiddleneck, coast
Fleabane, dwarf
Fleabane, hairy
Flixweed
Florida betony
Fumitory
Garlic, wild
Gera-
(cont'd)

[1]will be controlled or suppressed by this product when applied before weeds germinate

nium, Carolina[1]
Goldenrod, Canada
Goldenrod, Missouri
Goldenweed, common
Goosefoot, nettleleaf
Ground ivy (creeping Charlie)
Groundsel, common
Hawkweed
Henbane, black
Henbit[1]
Horsenettle, Carolina
Ironweed
Ivy, ground
Jacobs-ladder
Jimsonweed
Knapweed, black
Knapweed, diffuse
Knapweed, Russian
Knapweed, spotted
Knawel
Knotweed, common[1]
Knotweed, prostrate[1]
Kochia
Lambsquarters
Lambsthumb
Lespedeza, common[1]
Lettuce miners
Lettuce, prickly
Loosestrife, loosed-leaf
Mallow, common
Mallow, roundleaf
Mallow, Venice
Matchweed
Mayweed
Milkweed, climbing
Milkweed, common
Milkweed, honeyvine
Milkweed, Jerusalem whorled
Milkweed, western artichoke
Morningglory, ivyleaf
Morninglory, tall
Mustard, black[1]
Mustard, blue[1]
Mustard, Indian[1]
Mustard, tansy[1]
Mustard, treacle[1]
Mustard, tumble[1]
Mustard, wild[1]
Mustard, yellowtops[1]
Nettle, stinging
Nightshade, black
Oats wild
Onion, wild
Oxalis[1]
Parsley-piert[1]
Pigweed species[1]
Pineapple weed
Plantain, bracted
Plantain, broadleaf
Plantain, buckhorn
Poison ivy
Poison oak
Pokeweed
Purslane, common[1]
Pusley, Florida[1]
Radish, wild
Ragweed, common
Ragweed, giant (buffaloweed)
Ragweed, lance-leaf
Redmaids
Rocket, London[1]
Sheep sorrel
Shepherd's purse[1]
Smartweed, green
Smartweed, Pennsylvania
Sorrel, red
Sorrel, sheep
Sowthistle, annual
Speedwell, corn[1]
Spikeweed, common
Spotted catsear[1]
Spurge, leafy[1]
Spurge, spotted[1]
Spurweed
Storksbill, long
Sunflower, common
Sunflower, volunteer
Thistle
Thistle, bull
Thistle, Canada
Thistle, milk
Thistle, musk
Thistle, plumeless
Thistle, Russian
Velvetleaf
Vetch, common
Virginia buttonweed
Virginia pepperweed
Waterhemp
Wild violet
Willowherb
Woodsorrel, creeping[1]
Woodsorrel, yellow[1]
Wormwood
Yankeeweed
Yarrow common

[1]will be controlled or suppressed by this product when applied before weeds germinate

KILLS GRASSY WEEDS

Barley
Barnyard grass
Bluegrass, Annual (Poa annua)
Crabgrass, large
Crabgrass, smooth
Crabgrass, Southern
Foxtail
Foxtail yellow
Goldenrod
Goosegrass
Kikuyugrass (suppression)
Kyllinga, green
Marestail (horseweed)
Nutsedge, purple
Nutsedge, yellow
Ryegrass, annual and perennial
Sandbur (grass spur)
Smut grass

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING

If Empty:

- Non-refillable container. Do not refill or reuse this container.
- Place in trash or offer for recycling if available.

If Partly Filled:

- Call your local solid waste agency for disposal instructions.
- Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

IF IN EYES • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.

NOTE: When calling a poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call 24/7 toll-free 1-877-229-3763 for medical emergency information.**

Active Ingredients: 0.44% Dicamba (CAS # 1918-00-9), 0.10% Penoxsulam (CAS# 219714-96-2), and 1.10% Dithiopyr (CAS # 97886-45-8)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- After using this product wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

• This product is toxic to fish and highly toxic to aquatic invertebrates including oysters and shrimp. • To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. • Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. • Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Ground Water Advisory: This product has properties and characteristics associated with chemicals detected in groundwater. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. May result in ground water contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Non-Target Organism Advisory : This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DATOS RÁPIDOS

ACCIÓN DE DOS VÍAS

Mata y previene las malezas de la lista.

MATA POR CONTACTO

Usar con confianza en la mayoría de los tipos de césped, incluyendo el césped azul de Kentucky, el césped azul perenne, festuca, el raigrás perenne, el césped Bermuda, el césped ciempiés y el césped San Agustín (incluyendo floratam).
No deje nada liberado a las conjeturas en el control de malezas en el césped. Para tener césped hermoso libre de malezas.

COBERTURA

Cubre 4000 pies cuadrados de césped.
Si tiene preguntas o comentarios, llame gratis al 1-877-229-3724 o visite nuestro sitio web en www.BioAdvanced.com

INSTRUCCIONES DE USO Constituye una violación a la ley federal usar este producto de manera diferente a lo establecido en esta etiqueta.

ANTES DE USAR

Leer y seguir las instrucciones de la etiqueta.

RESTRICCIONES

- No aplicar este producto de modo que entre en contacto con personas o mascotas ya sea directamente o por goteo.
- Mantener a las personas y mascotas fuera del área durante su aplicación y hasta que el producto se haya secado.
- No permitir que las personas y las mascotas vuelvan a ingresar al área tratada hasta que el rocío se haya secado.
- No usar en césped que se usa para la venta u otro uso comercial como pasto, o para producción comercial de semillas o con fines investigativos.
- No aplicar directamente, o permitir que el rocío escurra en flores, arbustos ornamentales, árboles, u otras plantas que desee conservar, ya que puede ocurrir una lesión.
- No aplicar dentro de 10 pies de plantas sensibles, incluyendo las huertas de verduras.
- Los restos de los cortes deben dejarse en las áreas tratadas. No usar los restos de cortes de las áreas tratadas como mantillo o compost alrededor de flores, plantas decorativas, árboles o en jardines de vegetales.
- No tratar en céspedes que estén bajo estrés por sequía, inundación, fertilizador excesivo o suelos con sal, lesión por insectos, nematodos o enfermedad.
- No permitir que el producto ingrese en algún drenaje durante o después de la aplicación.
- No aplicar si se espera lluvia dentro de las 24 horas de la aplicación (para control mínimo 1 hora antes de que el producto haya tenido suficiente tiempo para secarse)

(continuación)

- No realizar aplicaciones si llueve.
- No aplicar si hay viento que podría hacer que el producto escurra en plantas sensibles.
- No rociar el producto en lagunas con peces, pozos, arroyos o lagos.
- No aplicar más de 1.5 lb ai/A de dithiopyr por año (cuando se usa producto adicional que contenga dithiopyr).
- No aplicar más de 8 libras de este producto por cada 1,000 pies cuadrados por aplicación (equivalente a 0.25 lb dithiopyr, 0.10 lb de dicamba, y 0.02 lb de penoxsulam por acre).
- No aplicar más de 32 onzas líquidas de este producto por cada 1,000 pies cuadrados por año (equivalente a 1 lb de dithiopyr, 0.40 lb de dicamba, y 0.08 lb de penoxsulam por acre).
- En el estado de New York no exceda 8 onzas líquidas del producto por cada 1,000 pies cuadrado por año (equivalente a 0.25 lb de dithiopyr por acre por año)

PRECAUCIONES

- Usar este producto solamente en céspedes establecidos.
- La lluvia excesiva dentro de las 24 horas luego de la aplicación
- Aplicar cuando el aire está calmo para evitar que se esparza en plantas no deseadas.

CUÁNDO USAR

Aplicar en cualquier momento desde la primavera hasta el otoño a hierbas de césped en crecimiento cuando las temperaturas estén entre 50-90 grados F para matar a las malezas de hoja ancha. Este producto no quemará el césped cuando la temperatura ascienda a 90 grados F.

DÓNDE USAR

EN CÉSPEDES ESTABLECIDOS

Céspedes de temporada fría: césped agrostis, pasto Kentucky, festuca (chewing, festuca roja trepadora, alta) y el riegrás perenne

Céspedes de temporada cálida: césped Bermuda (comunes o híbridos), festuca alta, kikuyugrass y césped zoysia, césped ciempiés, césped San Agustín, césped búfalo.

NO USAR EN NINGÚN OTRO TIPO DE CÉSPED incluyendo: dichondra, pasto chato, césped Bahía o hierba de césped que contenga tréboles que dese conservar.

CUÁNTO USAR

Usar como tratamiento por esparcimiento (tratamiento de todas las áreas del césped). Esta botella sirve para tratar hasta 4,000 pies cuadrados. Medir el área a rociar. Calcular los pies cuadrados multiplicando la longitud por el ancho. La tasa de aplicación máxima de este producto es 8 onzas líquidas por cada 1,000 pies cuadrados por aplicación. No aplicar más de 4 veces por año (excepto en New York). Consultar las restricciones de arriba para acceder a más información sobre las tasas máximas de uso.

CÓMO USAR

Es necesario cubrir todo el césped y las malezas para matar las malezas.

CÓMO COLOCAR Y ROCIAR

Cuando se use por primera vez, quitar el bloqueo de plástico de envío.



CONECTAR
1. Agitar bien antes de usar.
2. Conectar el rociador de manera segura a la manguera del jardín.
3. Asegúrese que el Rociador esté en posición cerrado OFF.
4. Abrir el grifo de agua.

ROCIAR
1. Extender la manguera hasta el área más lejana, apuntar la boquilla del rociador en dirección hacia donde desea rociar.
2. Girar el rociador a la posición ON.
3. Caminar a un ritmo constante mientras rocía, usando un movimiento oscilante y volver a regar la misma área tratada solo un poco. Caminar hacia atrás hacia el grifo para evitar caminar sobre el césped húmedo y tratado.

TERMINAR
1. Para finalizar el rocío, llevar el rociador hasta la posición OFF.
2. Apagar el grifo de agua.
3. Liberar la presión de agua en la manguera girando la válvula del rociador nuevamente a la posición ON hasta que el agua disminuya lentamente a un goteo, luego volver a girar la válvula del rociador a la posición OFF.

FRECUENCIA

Este producto puede aplicarse no antes de 35 días después de la aplicación anterior.

CÓMO FUNCIONA

Este producto contiene tres herbicidas selectivos – mata de manera selectiva malezas en el césped pero no dañará al césped.

PARA MEJORES RESULTADOS

CUÁNDO APLICAR
Rociar cuando estén presentes las malezas. • Temperaturas entre 50 - 90° F
• Do not apply while grass is emerging from dormancy. • No aplicar cuando el césped está emergiendo del letargo.

CORTADO No cortar durante 1 a 2 días después de haber rociado. • Los restos de los cortes deben dejarse en las áreas tratadas. No usar los restos de cortes de las áreas tratadas como mantillo o compost alrededor de flores, plantas decorativas, árboles o en jardines de vegetales.

REGADO Si el césped está seco, riegue 1 a 2 días antes de usar este producto.

CÓMO FUNCIONA ESTE PRODUCTO

Este producto es una mezcla probada de dos exterminadores de malezas luego de su aparición que mata las malezas actuales, más las que están por nacer para evitar nuevas malezas. Y no va a dañar a su césped, cuando se usa como se indica.

Dicamba – Dicamba es un herbicida selectivo que se usa para controlar un amplio espectro de malezas de hojas anchas. Tiene una ventana de aplicación amplia y ayuda en la quema de las malezas. Dicamba se absorbe rápidamente a través de las hojas de las plantas, los tallos, las raíces y funciona rápidamente alterando el crecimiento de la maleza. La tasa máxima de una sola aplicación es 0.10 lb ai/a.
Penoxsulam – Penoxsulam es un herbicida sistémico que se absorbe a través de las hojas y las raíces de las plantas. Las malezas tratadas con penoxsulam dejarán de crecer casi inmediatamente y se morirán a las 2 a 4 semanas luego de su aplicación. La tasa máxima de una sola aplicación es 0.02 lb ai/a.
Dithiopyr – Dithiopyr es un herbicida previo a la aparición que impide que las malezas de hoja ancha y las de hierba germinen. Tiene control post emergente de zácate cangrejo durante hasta 4 semanas después de que el zácate cangrejo germine y aparezca. Si el zácate cangrejo comienza a esparcirse, este producto no será efectivo y no debe aplicarse después de 4 semanas posteriores a la germinación. La tasa máxima de una sola aplicación es 0.25 lb ai/a.
Al igual que todos los exterminadores de malezas, la velocidad de acción depende de muchas condiciones (p. ej. tipo de maleza, temperatura) en un momento dado.

MATA MALEZAS DE HOJA ANCHA

Alkanet
Amaranto, palmera
Amaranto powell
Amaranto, espinoso
Aster, salvaje
Aster, con espinas
Aster, de brezo blanco
Begarweed, trepadora
Begarweed, enana
Betonia, común
Bindweed, de campo
Bindweed, erísimo[1]
Mielga negra[1]
Broomweed, campo
Buckwheat, tartary, salvaje,
trébol de carretilla, California
Silene Galio del bosque, común
Erechtites, Americana
Botón de oro, maíz
botón de oro, trepadora
Botón de oro, peluda
Botón de oro, cuerno rojo
Botón de oro, maleza dura
Botón de oro, del oeste común
Axonopus[1]
zanahoria salvaje
Ipomoea hederacea
Ipomoea hoja de hiedra
Ápio, silvestre
Manzanilla, maíz
Perejil, bur
Oreja de ratón (común)[1]
Oreja de ratón (oreja de raón)[1]
Trébol
Trébol, holandés
Trébol lonicera
Trébol, rojo
Trébol, blanco
Berberecho, maíz, vaca, blanca
Bardana menor común
Chícharo (común achicoria)
Copperleaf, hophornbeam
Cornflower (bachelor botón)
Crotón, trópico
Crotón, lanígero
Peludilla, púrpura
Lengua de vaca
Dalia japonesa
Margarita inglesa
Diente de león común
Diente de león falso
Maleza del dólar (pennywort)
Dragonhead, Americano
Eclipta
Prímulas de la noche
Semilla de lino falsa, semilla pequeña
Ortiguilla de la costa
Fleabane enana
Fleabane, peluda
Hierba de la sabiduría
Betonia de Florida
Fumitory
Ajo, silvestre
Geranio, Carolina[1]
Goldenrod, Canadá
Goldenrod, de Missouri
Goldenweed, común
Goosefoot, hoja de ortiga
Glechoma hederacea
Senecio, común
Vellosilla
Beleño, negro
Henbit[1]

Solano
(continuación)

[1]este producto lo controlará o suprimirá cuando se aplique antes de que las semillas germinen

americano, Carolina
Hiedra, del suelo
Jacobs-ladder
Jimsonweed
Centaurea, negra
Centaurea, difusa
Centaurea, rusa
Centaurea, manchada
Escleranto
Fallopia, común[1]
Sanguinaria mayor[1]
Kochia
Parra virgen, Quínoa blanca común
Lambsthumb
Lespedeza, común[1]
Lechuga minera
Lechuga espinosa
Arroyuela, de hoja suelta
Malva, común
Malva, de hoja redonda
Malva, Venecia
Matchweed
Mayweed
Algodoncillo de flor de sangre, trepadora
Algodoncillo de flor de sangre, común
Algodoncillo de flor de sangre, Jerusalén
Algodoncillo de flor de sangre, común,
Algodoncillo de flor de sangre, del oeste
Mostaza, negra[1]
Mostazas, azul[1]
Mostaza, India[1]
Mostaza, Tanaceto[1]
Mostaza, treacle[1]
Mostaza, tumble[1]
Mostaza, silvestre[1]
Mostaza, de centro amarillo[1]
Ortiga, con escozor
Solanácea, negra
Avena, silvestre
Cebolla salvaje
Oxalis[1]
Parsley-piert[1]
Especies de amaranto[1]
Matricaria discoidea
Plantago aristata
Plátano, hoja ancha
Plátano, buckhorn
Hiedra venenose
Roble venenoso
Fitolaca
Verdolaga, común[1]
Pusley, Florida[1]
Rabanito, silvestre
Ambrosía, común
Ambrosía, gigante (hierba búfalo)
Ambrosía, Calandrinia ciliata
Cohete, Londres[1]
Sheep sorrel
Bolsa de pastor[1]
Smartweed,
Smartweed verde,
Pasto de Pennsylvania,
Pasto rojo,
persicaria, cerraja,
Verónica anual, maíz[1]
Hierba espinosa, común
Espuela de gato[1]
Euphorbia, con hojas[1]
Euphorbia, manchada[1]
Espuela
Erodium, larga
Girasol, común
Girasol, voluntaria
Cardo
Cardo, bull
Cardo, Canadá
leche Cardo
musk Card,
Carduus acanthoides
Rusa
Hoja de terciopelo
Veza, común
Malva arbórea de Virginia
Lepidio Virginia
Amaranto de frutos ásperos
Violeta silvestre
Epilobium
Ocas, trepadora[1]
Ocas, amarilla[1]
Wormwood
Yankeeweed
Milenrama

[1]este producto lo controlará o suprimirá cuando se aplique antes de que las semillas germinen

MATA MALEZAS DEL CÉSPED

Cebada
Césped Barnyard
Césped azul anual
Zácate cangrejo, grande
Zácate cangrejo, suave
Zácate cangrejo, sur
Colo de zorro
Cola de zorro amarilla
Solidago
Goosegrass
Kikuyugrass (supresión)
Kyllinga, verde
Marestail (hierba carnicera)
Juncia real amarilla
Juncia real amarilla
Raigrás, anual y perenne
Cenchrus (brote de hierba)
Smut grass

ALMACENAMIENTO Y ELIMINACIÓN

No contaminar alimentos, agua o alimentos para mascotas por almacenamiento o eliminación:

ALMACENAMIENTO DEL PESTICIDA

Almacene en el contenedor original en un lugar seco y fresco, fuera del alcance de los niños, preferentemente en un gabinete de almacenamiento cerrado. Proteger de congelamiento.

ELIMINACIÓN DEL PESTICIDA Y MANIPULACIÓN DEL CONTENEDOR

Si está vacío:

- Contenedor no rellenable. No rellenar o volver a usar este contenedor.
- Colocar en la basura u ofrecer para su reciclado si fuese

If Partly Filled:

- llamar a la agencia de eliminación de desechos sólidos para conocer las posiblle
- Nunca colocar el producto sin usar en un drenaje de salida o de entrada.

AVISO: Las investigaciones y pruebas han determinado que las "Indicaciones de uso" corresponden al uso adecuado de este producto en condiciones esperadas. Hasta el punto en que aplica la ley, el comprador asume la responsabilidad de falta de rendimiento o seguridad si no se usa de acuerdo con las instrucciones.
Garantizamos la devolución del dinero: Si no está satisfecho con este producto, le devolveremos el dinero de la compra original.

PRIMEROS AUXILIOS

SI LO TRAGA • Llamar al centro de control de envenenamiento o al médico inmediatamente para asesoramiento sobre el tratamiento. • Hacer que la persona sorba un vaso de agua si puede tragar. • No inducir el vómito a menos que se lo indique un médico o el centro de control de envenenamiento. • No dar nada a una persona inconsciente.
EN CASO DE CONTACTO CON LOS OJOS • Mantener los ojos abiertos y enjuagar lenta y cuidadosamente con agua durante 15 -20 minutos. • Quitar los lentes de contacto, si los tuviera, luego de los primeros 5 minutos, continuar lavando. • Llamar al centro de control de envenenamiento o al médico para asesoramiento sobre el tratamiento.
SI ENTRA EN CONTACTO CON PIEL O ROPA • Quitar la ropa contaminada. • Lavar la piel inmediatamente con mucha agua durante 15-20 minutos. • Llamar al centro de control de envenenamiento o al médico para asesoramiento sobre el tratamiento.
NOTA: Cuando llame al centro de control de envenenamiento, lleve la etiqueta de este producto. Si acude en busca de tratamiento a una sala de emergencia o al consultorio de un médico, lleve esta etiqueta del producto para mostrarle al personal médico. **Puede llamar gratuitamente 24/7 al 1-877-229-3763 para** información sobre atención médica de urgencia.

DECLARACIONES DE PRECAUCIÓN

Peligroso para animales domésticos y humanos

PRECAUCIÓN

- Peligroso si lo traga.
- Causa irritación moderada en los ojos.
- Evitar contacto con los ojos, piel o ropa.
- Después de usar este producto, lavar cuidadosamente con agua y jabón luego de manipularlo y antes de comer, beber, masticar goma de mascar, usar tabaco o ir al baño

PELIGROS AMBIENTALES

• Este producto es tóxico para los el medio ambiente, no permita que el pesticida ingrese o se escurra por drenajes de tormenta, peces y altamente tóxico para otros invertebrados acuáticos incluyendo las ostras y los camarones. • Para proteger zanjas, alcantarillas o aguas superficiales. • Aplicar este producto en clima calmo, cuando no haya pronóstico de lluvia en las siguientes 24 horas ayudará a garantizar que el viento o la lluvia no dispersen o escurran el pesticida pesticida fuera del área de tratamiento. • Enjuagar el equipo usado para la aplicación en el área tratada ayudará a evitar que el producto ingrese en cuerpos de agua o sistemas de drenaje.

Asesoramiento sobre aguas subterráneas: Este producto tiene propiedades y características asociadas con químicos detectados en aguas subterráneas. Este producto puede escurrirse a las aguas subterráneas si se usa en suelos donde sea permeable especialmente donde la mesa de agua sea poco profunda. Puede resultar en contaminación de las aguas subterráneas. Su aplicación alrededor de una cisterna o pozo puede resultar en contaminación del agua potable o de las aguas subterráneas.

Asesoramiento sobre organismos que no forman parte del objetivo: Este producto es tóxico para plantas y puede tener impacto adverso en el forraje y en el hábitat de organismos que no sean parte de su objetivo, incluyendo los polinizadores, en áreas adyacentes al lugar del tratamiento. Proteja el forraje y el hábitat de los organismos que no son su objetivo siguiendo las instrucciones de la etiqueta que tienen el objetivo de minimizar la deriva del producto