FAST-MAX
MULTISYMPTOM RELIEF
3025204
POWERFUL NIGHTTIME RELIEF
3158844
DAY & NIGHT
COUGH & COLD RELIEF
MAXIMUM STRENGTH
NDC 63824-117-66
Mucinex FAST-MAX
SEVERE CONGESTION & COUGH
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant
Mucinex NIGHTSHIFT
COLD & FLU
Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Triprolidine HCl – Antihistamine
NIGHTTIME RELIEF FOR A BETTER MORNING
COUGH • FEVER • SORE THROAT
RUNNY NOSE • SNEEZING
✓ Controls Cough
✓ Relieves Nasal & Chest Congestion
✓ Thins & Loosens Mucus
FOR AGES 12+
TWO – 6 FL OZ (180 mL) bottles
TOTAL – 12 FL OZ (360 mL)

#1
Pharmacist Recommended Brand
2023-2024
Pharmacy Times
Celebrating 126 Years!
#1 COUGH & COLD BRAND DOCTORS TRUST
*IQVIA, ProVoice Survey, Dec. 2022
*U.S. News & World Report/Pharmacy Times 2023-2024


MAXIMUM STRENGTH
Drug Facts
Active ingredients (in each 20 mL) Purposes
Mucinex Fast-Max Severe Congestion & Cough
Dextromethorphan HBr 20 mg ..... Cough suppressant
Guaifenesin 400 mg ..... Expectorant
Phenylephrine HCl 10 mg ..... Nasal decongestant
Mucinex NIGHTSHIFT Cold & Flu
Acetaminophen 650 mg ..... Pain reliever/fever reducer
Dextromethorphan HBr 20 mg ..... Cough suppressant
Triprolidine HCl 2.5 mg ..... Antihistamine
Uses MUCINEX FAST-MAX SEVERE CONGESTION & COUGH
■ temporarily relieves these common cold and flu symptoms: ■ cough ■ minor aches and pains ■ sore throat ■ nasal congestion ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes due to hay fever
■ temporarily reduces fever
■ controls cough to help you get to sleep
Warnings
Liver warning (Nighttime Cold & Flu only): This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product
Allergy alert (Nighttime Cold & Flu only): Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning (Nighttime Cold & Flu only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. (Nighttime Cold & Flu only)
■ if you have ever had an allergic reaction to any other cough-cold or allergy drug containing acetaminophen
■ if you have (Nighttime Cold & Flu only) liver disease (Fast-Max Severe Congestion & Cough only)
■ heart disease (Fast-Max Severe Congestion & Cough only)
■ high blood pressure (Fast-Max Severe Congestion & Cough only)
■ diabetes (Fast-Max Severe Congestion & Cough only)
■ glaucoma (Nighttime Cold & Flu only)
■ trouble urinating due to an enlarged prostate gland
■ a breathing problem such as emphysema or chronic bronchitis
■ taking MAOI drugs
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ a cough that occurs with too much phlegm (mucus)
Ask a doctor before use if you have
■ liver disease (Nighttime Cold & Flu only)
■ heart disease (Fast-Max Severe Congestion & Cough only)
■ high blood pressure (Fast-Max Severe Congestion & Cough only)
■ diabetes (Fast-Max Severe Congestion & Cough only)
■ glaucoma (Nighttime Cold & Flu only)
■ trouble urinating due to an enlarged prostate gland
■ a breathing problem such as emphysema or chronic bronchitis
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ a cough that occurs with too much phlegm (mucus)
Drug Facts (continued)
Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin (Nighttime Cold & Flu only)
■ taking sedatives or tranquilizers (Nighttime Cold & Flu only)
When using this product
■ do not use more than directed
■ excitability may occur, especially in children (Nighttime Cold & Flu only)
■ marked drowsiness may occur (Nighttime Cold & Flu only)
■ alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime Cold & Flu only)
■ avoid alcoholic drinks (Nighttime Cold & Flu only)
■ be cautious when driving a motor vehicle or operating machinery (Nighttime Cold & Flu only)
Stop use and ask a doctor if
■ sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting (Nighttime Cold & Flu only)
■ cough gets worse or lasts more than 7 days (Nighttime Cold & Flu only)
■ fever gets worse or lasts more than 3 days (Nighttime Cold & Flu only)
■ nervousness, dizziness, or sleeplessness occur (Fast-Max Severe Congestion & Cough only)
■ redness or swelling is present (Nighttime Cold & Flu only)
■ new symptoms occur (Nighttime Cold & Flu only)
■ cough comes back or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. (Nighttime Cold & Flu only)
■ symptoms do not get better within 7 days or occur with fever (Fast-Max Severe Congestion & Cough only)
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. (Fast-Max Severe Congestion & Cough only)
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Overdose warning (Nighttime Cold & Flu only): Taking more than the recommended dose (overdose) may cause liver damage. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
MUCINEX FAST-MAX SEVERE CONGESTION & COUGH
■ do not take more than 6 doses in any 24-hour period
■ measure only with dosing cup provided
■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor
■ adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
■ children under 12 years of age: do not use
MUCINEX NIGHTSHIFT COLD & FLU
■ do not take more than directed (see Overdose warning)
■ do not take more than 4 doses in any 24-hour period
■ measure only with dosing cup provided
■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor
■ adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
■ children under 12 years of age: do not use
12 Total FL Oz

MAXIMUM STRENGTH COUGH & COLD BUNDLE
COUGH
FEVER
CHEST CONGESTION
SORE THROAT
SNEEZING
NASAL CONGESTION
RUNNY NOSE
MUCINEX FAST-MAX
SEVERE CONGESTION & COUGH
MUCINEX NIGHTSHIFT
COLD & FLU

AROUND THE CLOCK SYMPTOM RELIEF
MAXIMUM STRENGTH
Mucinex FAST-MAX
SEVERE CONGESTION & COUGH
Controls Cough
Relieves Nasal & Chest Congestion
Thins & Loosens Mucus
6 FL OZ (180 mL)
Mucinex NIGHTSHIFT
NIGHTTIME RELIEF FOR A BETTER MORNING

MAXIMUM STRENGTH
NDC 63824-540-66
Mucinex FAST-MAX
SEVERE CONGESTION & COUGH
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant
Controls Cough
Relieves Nasal & Chest Congestion
Thins & Loosens Mucus
6 FL OZ (180 mL)
FOR AGES 12+
DAYTIME RELIEF TO POWER THROUGH YOUR DAY

NIGHTTIME RELIEF FOR A BETTER MORNING
Mucinex NIGHTSHIFT
COLD & FLU
NIGHTTIME RELIEF FOR A BETTER MORNING
COUGH • FEVER • SORE THROAT
RUNNY NOSE • SNEEZING

FSA & HSA ELIGIBLE
SAVE UP TO 30%*
WHEN YOU PAY WITH YOUR FSA/HSA CARD!
1. Add Your FSA/HSA Card To Your Account
2. Shop FSA/HSA Eligible Products
3. Use Your FSA/HSA card at checkout
COVER YOUR OUT-OF-POCKET EXPENSES WITH PRE-TAX DOLLARS
*Plan details may vary, so save your receipts and speak with your health plan or benefits provider about the OTCs and personal healthcare products that are eligible in your plan.

YOU'VE GOT SYMPTOMS. WE'VE GOT RELIEF.
MucinexDM
Mucinex INSTA-SOOTHE
SORE THROAT + SOOTHING COMFORT
3-in-1 FORMULA
Mucinex FAST-MAX
NIGHT-TIME SEVERE COLD & FLU
Mucinex KICKSTART SEVERE COLD & FLU
ALL-IN-ONE
USE AS DIRECTED

FAST-MAX
MULTISYMPTOM RELIEF
3025204
POWERFUL NIGHTTIME RELIEF
3158844
DAY & NIGHT
COUGH & COLD RELIEF
MAXIMUM STRENGTH
NDC 63824-117-66
Mucinex FAST-MAX
SEVERE CONGESTION & COUGH
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant
Controls Cough
Relieves Nasal & Chest Congestion
Thins & Loosens Mucus
Mucinex NIGHTSHIFT
COLD & FLU
Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Triprolidine HCl – Antihistamine
NIGHTTIME RELIEF FOR A BETTER MORNING
COUGH • FEVER • SORE THROAT
RUNNY NOSE • SNEEZING
FOR AGES 12+
TWO – 6 FL OZ (180 mL) bottles
TOTAL – 12 FL OZ (360 mL)
FOR AGES 12+

Drug Facts (continued)
Other information
Inactive ingredients (Mucinex Fast-Max Severe Congestion & Cough)
Inactive ingredients (Mucinex Nightshift Cold & Flu)
Stored take Mucinex Fast-Max Severe Congestion & Cough and Mucinex Nightshift Cold & Flu at the same time. Always wash your hands before and after each dose of Mucinex liquid.
TAKE ONLY AS DIRECTED.

Drug Facts (continued)
Active ingredients (in each 20 mL) Purposes
Mucinex Fast-Max Severe Congestion & Cough
Mucinex Nightshift Cold & Flu

Drug Facts (continued)
Uses
Warnings
Directions
Other information
Inactive ingredients