EPA Reg. No. 92564-17 EPA Est. No indicated by 3rd and 4th digits of the batch number on this package. (65) = 92564-TX-1 (57) = (07)106-GA-003 (46) = 88746-NIO-1
Product of India. Distributed by: SBM Life Science Corp. PO Box 325, Morrisville, NC 27560, BioAdvanced® is a registered trademark of SBM Life Science Corp.

BIOADVANCED® SCIENCE-BASED SOLUTIONS TOMATO & VEGETABLE INSECT KILLER READY-TO-SPRAY

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
IF INHALED • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel.
You may call toll-free (877) 229-3763 for medical emergency information.
Active ingredient: 0.75% Cyfluthrin [CAS # 68359-37-5].

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

• Harmful if swallowed or inhaled. • Causes eye irritation. • Avoid contact with skin, eyes, or clothing. • Wash thoroughly with soap and water after handling. • Do not use kitchen utensils for measuring. Keep utensils used for measuring with product and away from children.

ENVIRONMENTAL HAZARDS

• This pesticide is toxic to fish and other aquatic organisms. • This pesticide is highly toxic to bees exposed to direct treatment or residues on plants or weeds. Do not apply this product or allow it to drift onto plants or weeds on which bees are actively foraging. • To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

PRESS TO RESEAL

501437193 20092511

BIOADVANCED®
SCIENCE-BASED SOLUTIONS

TOMATO & VEGETABLE

INSECT KILLER

READY-TO-SPRAY

Just one application quickly kills listed insects

INSECTS CONTROLLED

APHIDS | ADULT JAPANESE BEETLES | CUTWORMS | TOMATO HORNWORMS

WHERE TO USE

LEAFY VEGETABLES | FRUITING VEGETABLES

ROOT VEGETABLES†† | CUCURBITS

††except sugarbeet

QUICK FACTS

RAINPROOF IN ONE HOUR | FAST ACTING-KILLS BY CONTACT

STOP-READ THIS ENTIRE LABEL BEFORE USE

Indicaciones en español después de las de inglés

FOR OUTDOOR RESIDENTIAL USE ONLY

8 40216 20085 2

FOR QUESTIONS OR COMMENTS?
Call toll-free
(877) 229-3724
or visit us online at
www.bioadvanced.com

SATISFACTION 100% GUARANTEED†

†Or your money back.

PEEL HERE

501437193 200925V1

QUICK FACTS

KILLS
Aphids, Caterpillars, Japanese Beetles (adults), Leaf-feeding beetles, Stink Bugs (including Brown Marmorated Stink Bugs), Whiteflies and other listed pests.

COVERAGE
Makes up to 64 gallons of spray

WHERE TO USE
Vegetables, Peanuts and Grapes

WHEN TO USE
When insects or insect damage first appear

HOW MUCH TO USE
No measuring needed. Dilutes automatically.

WHEN TO HARVEST
See plant specific restrictions for pre-harvest interval. If no pre-harvest interval is listed, the product may be applied up to and including day of harvest.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.

BEFORE YOU USE

Read and follow these directions when using.

- Apply the product to flowering plants during early morning or late evening, when bees are not present.
- Do not spray the product into fish pools, ponds, streams, or lakes.
- Remove animal feeding dishes prior to application. Keep children and pets away from treated areas until spray has dried.
- Do not water the treated area to the point of run-off.
- Do not make application during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
  (1) Treatment to soil or vegetation around structures;
  (2) Applications to lawns, turf, and other vegetation;
  (3) Applications to building foundations, up to a maximum height of 3 feet.
  Other than applications to building foundations, all outdoor applications to

(cont'd)

BEFORE YOU USE *(cont'd)*

impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack and crevice applications, only.

- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

WHEN TO USE

Spray when listed insects or damage first appear. Refer to specific use area for additional instructions.

HOW TO ATTACH AND SPRAY

- Shake well before using.
- Adjust nozzle on your sprayer to desired spray pattern.
- Hold sprayer about 12 inches from foliage.
- Spray leaves and tender new shoots.
- To kill listed insects that may hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray, but not dripping.

CONNECT



1. Shake well before use.
2. Securely Connect sprayer to garden hose.
3. Be sure that ON/OFF Switch is in OFF position, toward hose end of sprayer.
4. Turn on water at faucet.
5. For first time use, remove plastic shipping lock from beneath switch.

SPRAY

1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed.
2. Begin spraying - Push ON/OFF Switch forward into ON position.
3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap treated areas. Work back toward faucet to avoid walking over wet, treated grass.

FINISH

1. To stop spraying, push ON/OFF Switch to OFF position.
2. Turn off water at faucet.
3. Relieve water pressure in hose by turning ON/OFF Switch back to ON position until water slows to a drip, then return ON/OFF switch to OFF position.
4. Disconnect sprayer from hose.

FRUITING VEGETABLES

Eggplant, Groundcherry, Pepino, Pepper (includes: bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, and Tomato

PESTS

Beet armyworm (1st and 2nd instar), Cabbage looper, Celery leaftier, Colorado potato beetle, European corn borer, Flea beetles, Garden webworm, Leafminers, Pepper weevil, Potato aphid, Potato leafhopper, Southern armyworm (1st and 2nd instar), Stink bugs (including Brown Marmorated Stink Bugs), Tarnished plant bug, Thrips (except Thrips palmi), Tomato fruitworm (corn earworm), Tomato hornworm, Tomato pinworm, Variegated cutworm, Whitefly (adult), Western yellowstriped armyworm (1st and 2nd instar)

Restrictions:

- **Pre-harvest Interval:** Do not harvest until at least 7 days after application, except for tomato, which has no PHI.
- Do not spray more than 6 times per year.

BRASSICA (COLE) LEAFY VEGETABLES:

Broccoli, Broccoli raab (rapini), Chinese (gai lon) broccoli, Brussels sprouts, Cabbage, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, and Turnip greens.

PESTS

Alfalfa looper, Armyworm (1st and 2nd instar), Beet armyworm (1st and 2nd instar), Cabbage flea beetle, Cabbage looper, Cabbage webworm, Corn earworm, Cutworms, Diamondback moth (larvae), Fall armyworm (1st and 2nd instar), Grasshoppers, Imported cabbageworm, Japanese beetle (adult), Lygus bug, Meadow spittlebug, Potato leafhopper, Southern armyworm (1st and 2nd instar), Stink bugs, Southern cabbageworm, Tarnished plant bug, Thrips , Vegetable weevil (adult), Whitefly (adult), Yellowstriped armyworm (1st and 2nd instar)

Restrictions:

- Do not spray more than 4 times per year.

LEAFY VEGETABLES

Amaranth (Chinese spinach), Arugula (rouquette), Cardoon, Celery, Chinese celery, Celtuce, Chervil, Chrysanthemum (edible-leaved and garland), Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Endive (escarole), Florence fennel,

(cont'd)

LEAFY VEGETABLES *(cont'd)*

Lettuce (head and leaf), New Zealand spinach, Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, Swiss chard, Vine spinach

PESTS

Alfalfa looper, Beet armyworm (1st and 2nd instar), Cabbage looper, Corn earworm, Cutworms, Diamondback moth (larvae), European corn borer, Fall armyworm (1st and 2nd instar), Flea beetles, Grasshoppers, Green cloverworm, Imported cabbageworm, Japanese beetle (adult), Leafhoppers, Lygus bug, Meadow spittlebug, Saltmarsh caterpillar, Southern armyworm (1st and 2nd instar), Stink bugs, Tarnished plant bug, Thrips, Vegetable weevil (adult), Whitefly (adult), Yellowstriped armyworm (1st and 2nd instar)

Restrictions:

- Do not spray more than 4 times per year.

CUCURBITS (EXCEPT CROPS GROWN FOR SEED):

Balsam apple, Balsam pear, Bitter melon, Chayote, Chinese cucumber, Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes: hyotan, cucuzza, henchmia and Chinese okra), Muskmelon (includes: cantaloupe, true cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Summer squash (includes: crookneck squash, scallop squash, straightneck squash, vegetable marrow, and zucchini), Watermelon, Winter squash (includes: butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash)

PESTS

Armyworm (1st and 2nd instar), Cabbage looper, Corn earworm, Cutworms, Cucumber beetle, Grasshoppers, Lygus bug, Melonworm, Pickleworm, Potato leafhopper, Rindworm, Stink bugs, Stripped cucumber beetle, Tarnished plant bug, Tobacco budworm, Whitefly (adult)

Restrictions:

- Do not spray more than 4 times per year.

POTATO AND OTHER TUBEROUS AND CORM VEGETABLES:

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible canna, Cassava (bitter and sweet), Chayote root, Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, True yam, Turmeric, Yam bean

(cont'd)

POTATO AND OTHER TUBEROUS AND CORM VEGETABLES: *(cont'd)*

PESTS

Aphids, Cabbage looper, Colorado potato beetle, Cutworms, European corn borer, Flea beetles, Potato psyllid, Potato tuberworm, Sweetpotato weevil (adults), Tarnished plant bug

Restrictions:

- Do not spray more than 6 times per year.

ROOT VEGETABLES (EXCEPT SUGARBEET):

Garden beet, Edible burdock, Carrot, Celeriac, Turnip-rooted chervil, Chicory, Ginseng, Horseradish, Turnip-rooted parsley, Parsnip, Radish, Oriental radish, Rutabaga, Salsify (black, Spanish, and oyster plant), Skirret, Turnip

PESTS

Aster leafhopper, Carrot weevil, Cutworms, Flea beetles, Potato leafhopper

Restrictions:

- Do not spray more than 5 times per year.

SWEET CORN

PESTS

Armyworm (1st and 2nd instar), Bean leaf beetle, Black cutworm, Cereal leaf beetle, Chinch bug, Click beetle (adult), Corn earworm, Corn rootworms (adult), Corn silk fly, European corn borer, Fall armyworm (1st and 2nd instar), Flea beetle, Granulate cutworm, Grape colaspis (adult), Grasshoppers, Japanese beetle (adult), June beetle (adult), Leafhoppers, Masked chafer (adult), Sand hill cutworm, Southern armyworm (1st and 2nd instar), Southern corn leaf beetle, Southwestern corn borer, Stalk borer, Stink bugs, Webworm, Western bean cutworm, Yellowstriped armyworm (1st and 2nd instar)

Restrictions:

- Do not spray more than 10 times per season.

PEANUT

PESTS

Armyworm (1st and 2nd instar), Bean leaf beetle , Beet armyworm (1st and 2nd instar), Cutworms, Corn earworm, Corn rootworms (adult), Fall armyworm (1st and 2nd instar), Green cloverworm, Grape colaspis (adult), Grasshoppers,

(cont'd)

PEANUT *(cont'd)*

PESTS

Japanese beetle (adult), June beetle (adult), Potato leafhopper, Rednecked peanutworm, Stink bugs, Threecornered alfalfa hopper, Southern armyworm (1st and 2nd instar), Soybean looper, Thrips, Vegetable weevil, Velvetbean caterpillar, Whitefringed beetle (adult), Whitefly (adult)

Restrictions:

- **Pre-harvest Interval:** Do not harvest until at least 14 days after application.
- Do not spray more than 3 times per year

GRAPES

Includes: Table grape, Raisin, Wine and Muscadine grape

PESTS

Climbing cutworm, Grape berry moth, Grape bud beetle, Grape cane gallmaker (adult), Grape flea beetle, Grape leaffolder, Grape leafhopper, Grape leafroller, Grape mealybug (crawlers), Glassywinged sharpshooter, Grape leaf skeletonizer, Omnivorous leafroller, Orange tortrix, Thrips, Variegated leafhopper, Western grape leaf skeletonizer

Restrictions:

- **Pre-harvest Interval:** Do not harvest until at least 3 days after application.
- Do not spray more than 4 times per year.

FOR BEST RESULTS:

Do not water the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply.

This product is ready to spray. Just attach this bottle to your hose according to directions. Water automatically mixes with the concentrate as you spray.



People and pets must not reenter the treated area until after spray has dried.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

PESTICIDE DISPOSAL AND CONTAINER HANDLING

If Empty:

- Non-refillable container. Do not refill or reuse container.
- Place empty container in trash or offer for recycling if available.

If Partly Filled:

- Call your local solid waste agency for disposal instructions.
- Never place unused product down any indoor or outdoor drain.

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.
- Call a poison control center or doctor for treatment advice.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel.

You may call toll-free (877) 229-3763 for medical emergency information.
Active ingredient: 0.75% Cyfluthrin [CAS # 68359-37-5].

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed or inhaled.
- Causes eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling.
- Do not use kitchen utensils for measuring. Keep utensils used for measuring with product and away from children.

ENVIRONMENTAL HAZARDS

- This pesticide is toxic to fish and other aquatic organisms.
- This pesticide is highly toxic to bees exposed to direct treatment or residues on plants or weeds. Do not apply this product or allow it to drift onto plants or weeds on which bees are actively foraging.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DATOS RÁPIDOS

MATA
Áfidos, orugas, escarabajos japoneses (adultos), escarabajos comedores de hojas, chinches hediondas (incluso las chinches hediondas marmoradas marrones), moscas blancas y otras plagas.

COBERTURA
Rinde hasta 64 galones (240 litros)

DÓNDE USAR
Vegetales, Cacahuate y Uvas

CUÁNDO USAR
Cuando los insectos o el daño de los insectos aparecen por primera vez

CUÁNTO USAR
No es necesario medir. Se diluye automáticamente.

CUÁNDO COSECHAR
Consultar las restricciones específicas para obtener el intervalo previo a la cosecha. Si no se indica ningún intervalo previo a la cosecha, el producto se puede aplicar hasta incluso el día de la cosecha.

INDICACIONES DE USO
Es una violación de la Ley Federal utilizar este producto de manera contraria a la indicada en la etiqueta. Para obtener mejores resultados, leer y seguir todas las indicaciones de la etiqueta.

ANTES DE USAR

Leer y seguir las indicaciones de uso.

- Aplicar el producto a las plantas florecientes a la mañana temprano o a la última hora de la tarde, cuando las abejas no están presentes.
- No rociar el producto en estanques para peces, lagunas, arroyos o lagos.
- Retirar los platos para la alimentación de animales antes de la aplicación. Mantener a los niños y mascotas alejados de las áreas tratadas hasta que el rocío se haya secado.
- No regar el área tratada hasta el punto de escurrimiento.
- No realizar la aplicación durante la lluvia.
- Todas las aplicaciones en exteriores deben limitarse solamente a tratamientos localizados o en grietas y huecos, excepto para los siguientes usos permitidos:
  (1) Tratamiento al suelo o vegetación alrededor de estructuras;
  (2) Aplicaciones al césped, hierba y otra vegetación;
  (3) Aplicaciones a cimientos de edificios, hasta una altura máxima de 3 pies (90 cm).
  En aplicaciones distintas a la de cimientos de edificios, todas las aplicaciones en exteriores a superficies impermeables tales como aceras, entradas para vehículos, patios, galerías y superficies estructurales (tales como ventanas, puertas y aleros) están limitadas solamente a aplicaciones localizadas y de grietas y huecos.

(continúa)

ANTES DE USAR *(continúa)*

- Está prohibida la aplicación directamente en cloacas o canales de desagüe, o en cualquier área como una alcantarilla donde se produce el desagüe a cloacas, bocas de tormenta, cursos de agua o hábitat acuático. No permitir que el producto ingrese en algún desagüe durante o después de la aplicación.

CUÁNDO USAR

Rociar cuando aparece por primera vez el daño o los insectos de la lista. Consultar al área de uso específico para instrucciones adicionales.

CÓMO COLOCAR Y ROCIAR

- Agitar bien antes de usar.
- Ajustar la boquilla del rociador al patrón de rocío deseado.
- Sostener el rociador a aproximadamente 12 pulgadas (30 cm) del follaje.
- Rociar las hojas y los nuevos brotes tiernos.
- Para matar las plagas de la lista que se pueden esconder, rociar el lado inferior de las hojas. Aplicar hasta que las hojas estén cubiertas de forma pareja con el rocío, pero que no goteen.

CONECTAR



1. Agitar bien antes de usar.
2. Conectar el rociador de manera segura a la manguera del jardín.
3. Asegurarse que el interruptor ON/OFF esté en posición OFF, en el extremo de la manguera del rociador.
4. Abrir el grifo de agua.
5. Cuando se use por primera vez, quitar el bloqueo de plástico de envío de atrás del interruptor.

ROCIAR

1. Extender la manguera hasta el área más lejana, apuntar la boquilla del rociador en dirección hacia donde desea rociar.
2. Comenzar a rociar - Pulsar el interruptor ON/OFF hacia adelante a la posición ON.
3. Caminar a un ritmo constante mientras rocía, usando un movimiento oscilante y volver a regar la misma área tratada solo un poco. Caminar hacia atrás hacia el grifo para evitar caminar sobre el césped húmedo y tratado

TERMINAR

1. Para finalizar el rocío, pulsar el interruptor ON/OFF a la posición OFF.
2. Apagar el grifo de agua.
3. Liberar la presión de agua en la manguera pulsando el interruptor ON/OFF nuevamente a la posición ON hasta que el agua disminuya lentamente a un goteo, luego volver a pulsar el interruptor ON/OFF a la posición OFF.
4. Desconectar el rociador de la manguera.

HORTALIZAS QUE DAN FRUTOS

Berenjena, alquequenje, pepino, pimiento (incluye: ají, chile, pimiento para cocinar, pimentón y pimiento dulce), tomatillo y tomate.

PLAGAS

Gusano de la remolacha (1° y 2° estadio), oruga de la col, cigarrero del apio, escarabajo de la patata de Colorado, barrenador del maíz, escarabajo pulga, gusanos telerañeros de jardín, minadores de la hoja, oruga tejedora de jardín, minador de las hojas, gorgojo del chile, áfido de la papa, *(continúa)*

HORTALIZAS QUE DAN FRUTOS *(continúa)*

PLAGAS

saltarilla de la papa, gusano soldado de sur (1° y 2° estadio), chinche hedionda (incluyendo las chiches hediondas marrones marmóreas), chinche manchada de la planta, trips (excepto trips palmi),gusano del tomate (gusano de lamazorca del maíz), gusano del cuerno del tomate, minador del tomate, gusano cortador multicolor, mosca blanca (adulto), gusano soldado occidental de rayas amarillas (1° y 2° estadio).

Restricciones:

- **Intervalo previo a la cosecha:** no cosechar hasta al menos 7 días después de la aplicación, excepto los tomates que no tienen intervalo previo a la cosecha.
- No rociar más de 6 veces por año.

HORTALIZAS DE HOJA DEL GÉNERO BRASSICA (COL)

Brócoli, brócoli raab (rapini), brócoli chino (gai lon), repollito de Bruselas, col, col chino (bok choy), col chino (napa), col de mostaza china (gai choy), coliflor, brócoli de Cavalo, berzas, col rizada, colibano, mizuna, hojas de mostaza, espinaca de mostaza, colza y hojas de nabo.

PLAGAS

Oruga de la alfalfa, gusano soldado (1° y 2° estadio), gusano de la remolacha (1° y 2° estadio), escarabajo pulga de la col, oruga de la col, oruga tejedora de la col, gusano de la mazorca del maíz, gusano cortador, polilla dorso de diamante, gusano cogollero, (1° y 2° estadio), saltamontes, blanquita de la col, escarabajo japonés (adulto), chinche lygus, cigarra espumadora, saltarilla de la papa, gusano soldado del sur (1° y 2° estadio), chinche hedionda, oruga de la col del sur, chinche manchada de la plancha, trips, gorgojo de las hortalizas (adulto), mosca blanca (adulta), gusano soldado occidental de rayas amarillas (1° y 2° estadio).

Restricciones:

- No rociar más de 4 veces por año.

HORTALIZAS CON HOJAS

Amaranto (espinaca china), arúgula (rouquette), cardo, apio, apio chino, lechuga china, perifollo, crisantemo (con hojas comestibles y florigelio), colleja, berro (de jardín y tierras altas), diente de león, acedera (acetosa), endivia (escarola), hinojo de Florencia, lechuga (cabeza y hojas), espinaca de Nueva Zelanda, cenizo, perejil, verdolaga (de jardín e invierno), radicheta (achicoria roja), ruibarbo, espinaca, acelga, espinaca de enredadera.

PLAGAS

Oruga de la alfalfa, gusano de la remolacha (1° y 2° estadio), oruga de la col, gusano de la mazorca del maíz, gusano cortador, polilla dorso de diamante (larva), barrenador del maíz, gusano cogollero (1° y 2° estadio), escarabajo pulga, saltamontes, gusano verde del trébol, blanquita de la col, escarabajo japonés (adulto), saltarilla, chinche lygus, cigarra espumadora, gusano peludo, gusano soldado del sur (1° y 2° estadio), chinche hedionda, chinche manchada de la planta, trip, gorgojo de las hortalizas (adulto), mosca blanca (adulta), gusano soldado occidental de rayas amarillas (1° y 2° estadio).

Restricciones:

- No rociar más de 4 veces por año.

HORTALIZAS CUCURBITÁCEAS (EXCEPTO CULTIVOS PARA SEMILLA)

Balsamina, pera bálsamo, melón amargo, cayote, pepino chino, calabaza de cera (melón conservante de China), patilla, pepino, pepinillo, calabacino comestible (incluye: hyotan, cucuzza, hechima y quimbombó de China), melón reticulado (incluye: melón cantalupo, cantalupo, casaba, melón Crenshaw, melón Pershaw dorado, melón dulce, melones "Honey ball", melón mango, melón de Persia, melón piña, melón Santa Claus y melón serpiente), zapallo, calabaza de verano (incluye: calabaza de cuello curvo, calabaza escalopini, calabaza de cuello derecho, cucúrbita y zucchini), sandía, calabaza de invierno (incluye: calabaza de sidra, calabaza, calabaza Hubbard, cucúrbita pepo y calabaza spaghetti).

PLAGAS

Gusano soldado (1° y 2° estadio), oruga de la col, gusano de la mazorca del maíz, gusano cortador, escarabajo del pepino, saltamontes, chinche lygus, gusano del melón, gusano perforador del pepino, saltarilla de la papa, perforador de la vid de la calabaza, chinche hedionda, escarabajo con rayas del pepino, chinche manchada de la planta, gusano del tabaco, mosca blanca (adulta).

Restricciones:

- No rociar más de 4 veces por año.

PATATAS Y OTROS TUBÉRCULOS Y BULBOS

Arracacha, arrurruz, alcachofa (china y aguaturma), achira comestible, mandioca (dulce y amarga), raíz de cayote, chufa, colocasia esculenta (taro), jengibre, lerén, patata, batata, cambrón, ñame, cúrcuma, nupe.

PLAGAS

Áfidos, oruga de la col, escarabajo de la patata de Colorado, gusano cortador, barrenador del maíz, escarabajo pulga, psílido de la patata, gusano de los tubérculos, gorgojo de la patata (adulto), chinche manchada de la planta.

Restricciones:

- No rociar más de 6 veces por año.

HORTALIZAS CON RAÍCES (EXCEPTO LA REMOLACHA AZUCARERA)

Remolacha, bardana comestible, zanahoria, apio, perifollo enraizado al nabo, endivia, jengibre, rábano picante, perejil enraizado al nabo, chiviría, rábano, rábano oriental, colinabo, salsifí (negro, de España, Trapopogon porrifolius), Sium sisarum, nabo.

PLAGAS

Macrosteles quadrilineatus, gorgojo de la zanahoria, gusano cortador, escarabajo pulga, saltarilla de la papa.

Restricciones:

- No rociar más de 5 veces por temporada.

MAÍZ DULCE

PLAGAS

Gusano soldado (1° y 2° estadio), Cerotoma trifurcata, gusano cortador negro, escarabajo del cereal, chinche del maíz, escarabajo saltarín (adulto), gusano de la mazorca del maíz,

(continúa)

MAÍZ DULCE *(continúa)*

PLAGAS

gusano de la raíz del maíz (adulto), mosca de la mazorca, barrenador del maíz, gusano cogollero (1° y 2° estadio), escarabajo pulga, Cuncunilla granulosa, colaspis (adulto), saltamontes, escarabajo japonés (adulto), melotonta (adulto), saltarilla, escarabajo enmascarado (adulto), gusano cortador, gusano soldado del sur (1° y 2° estadio), escarabajo de la hoja del maíz del sur, barrenador del maíz del suroeste, gusano barrenador del tallo, chinche hedionda, oruga tejedora, gusano cortador occidental del frijol, gusano soldado de rayas amarillas (1° y 2° estadio).

Restricciones:

- No rociar más de 10 veces por año.

CACAHUATE

PLAGAS

Gusano soldado (1° y 2° estadio), Cerotoma trifurcata, gusano de la remolacha (1° y 2° estadio), gusano cortador, gusano de la mazorca del maíz, gusano de la raíz del maíz (adulto), gusano cogollero (1° y 2° estadio), gusano verde del trébol, colaspis (adulto), saltamontes, escarabajo japonés (adulto), melotonta (adulto), saltarilla de la papa, cogollero del maní, chinche hedionda, saltón de tres cuernos de la alfalfa, gusano soldado del sur (1° y 2° estadio), oruga de la soja, trips, gorgojo de las hortalizas, orugas de las leguminosas, gorgojo de la alfalfa (adulto), mosca blanca (adulta).

Restricciones:

- **Intervalo previo a la cosecha:** no cosechar hasta al menos 14 días después de la aplicación.
- No rociar más de 3 veces por año

UVAS

Incluye: uva de mesa, uva pasa, uva de vino y uva muscadina

PLAGAS

Gusano cortador trepador, polilla de la uva, escarabajo del brote de la uva, Coleoptera curculionidae (adulto), escarabajo pulga de la uva, taladrador de la uva, saltarilla de la uva, enrollador de la uva, cocos de la uva (etapa de arrastre), chicharrita de las alas cristalinas, descarnador de la hoja de la vid, gusano enrollador omnívoro, tortrix de la naranja, trips, saltarilla multicolor, descarnador de la vid.

Restricciones:

- **Intervalo previo a la cosecha:** no cosechar hasta al menos 3 días después de la aplicación.
- No rociar más de 4 veces por año

PARA OBTENER MEJORES RESULTADOS:

No regar el follaje dentro de la hora después de la aplicación. Para evitar daños graves a las plantas, eliminar las plagas a principios de la temporada antes de que se multipliquen.

Este producto está listo para pulverizar. Solo tiene que colocar la botella en la manguera de acuerdo con las instrucciones. El agua se mezcla automáticamente con el concentrado cuando pulverice.



Las personas y las mascotas pueden volver a ingresar al área tratada una vez que se haya secado.

NOTA: Investigaciones y pruebas han determinado que las "Indicaciones de uso" son las adecuadas para el uso apropiado de este producto bajo las condiciones previstas. El Comprador asume la responsabilidad, en la medida en que esté de acuerdo con la ley vigente, por la falta de rendimiento o seguridad si el producto no es utilizado según las indicaciones de uso.

Garantía de devolución de dinero: Si usted no está satisfecho con el producto, gustosamente le reembolsaremos el monto original del precio de compra.

ALMACENAMIENTO Y ELIMINACIÓN

ALMACENAMIENTO DEL PESTICIDA

Almacenar en el contenedor original y en un lugar fresco y seco, fuera del alcance de los niños, preferentemente en un armario bajo llave.

ELIMINACIÓN DEL PESTICIDA Y MANIPULACIÓN DEL CONTENEDOR

Si está vacío:

- Contenedor no reusable. No volver a rellenar ni utilizar este contenedor.
- Colocar el contenedor vacío en la basura o entregarlo para reciclar si fuera posible.

Si está parcialmente lleno:

- Llamar a la agencia local de residuos sólidos para solicitar indicaciones para la eliminación.
- Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.

PRIMEROS AUXILIOS

EN CASO DE INGESTIÓN

- Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.
- Hacer que la persona beba tragos de agua, si puede tragar.
- No inducir el vómito a menos que lo indique el médico o el centro de control de envenenamiento.
- No administrar nada por boca a una persona inconsciente.

EN CASO DE CONTACTO CON LOS OJOS

- Mantener los ojos abiertos y enjuagar en forma lenta y suave con agua durante 15 a 20 minutos.
- Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos, y luego continuar enjuagando los ojos.
- Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.

EN CASO DE INHALACIÓN

- Trasladar a la persona a un lugar fresco.
- Si la persona no respira, llamar al 911 o a una ambulancia, luego dar respiración artificial, preferentemente boca a boca, si fuera posible.
- Llamar a un médico o a un centro de control de envenenamiento para mayores consultas sobre el tratamiento.

(continúa)

PRIMEROS AUXILIOS *(continúa)*

NOTA: Al llamar a un centro de control de envenenamiento, tener la etiqueta del producto a mano. En caso de solicitar tratamiento en una sala de emergencia o consultorio médico, llevar la etiqueta de este producto para mostrarla al personal médico.
Puede llamar gratuitamente al 877-229-3763 para información sobre atención médica de urgencia.

Ingrediente activo: 0.75% ciflutrina [CAS Nº 68359-37-5].

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
DECLARACIONES DE PRECAUCIÓN

Riesgos para seres humanos y animales domésticos
PRECAUCIÓN

- Es perjudicial si se ingiere o inhala.
- Causa irritación en los ojos.
- Evitar el contacto con la piel, los ojos o la ropa.
- Lavarse prolijamente con agua y jabón después de manipular este producto.
- No utilizar utensilios de cocina para medir. Mantener los utensilios utilizados para medir con producto y alejados de los niños.

RIESGOS AMBIENTALES

- Este pesticida es tóxico para los peces y otros organismos acuáticos.
- Este pesticida es altamente tóxico para las abejas expuestas al tratamiento directo o residuos en plantas o malezas. No aplicar este producto o permitir que se disperse entre las plantas o malezas en los que las abejas buscan activamente.
- Para proteger el medio ambiente, no permitir que el pesticida ingrese o se escurra en las bocas de tormenta, zanjas, alcantarillas o aguas superficiales. Aplicar este producto cuando el clima esté calmo y no haya probabilidad de lluvia para las 24 horas siguientes ayudará a garantizar que el viento o la lluvia no dispersen o escurran el pesticida del área de tratamiento. Enjuagar el equipo de aplicación sobre el área tratada ayudará a evitar que ingrese en cursos de agua o los sistemas de desagüe.