NEUTERED YOUNG MALE



 A high protein diet helps maintain muscle mass, protein also yields less net energy than carbohydrates and L-carnitine helps transport fatty acids into the mitochondria for metabolism.



• Neutered male cats have a tendency to gain fat mass, leading to obesity which is a major risk factor for diabetes mellitus. This formula is reduced in caloric density, with a moderate carbohydrate content.





Available in 400g, 1.5kg, 3.5kg and 10kg bag

Approximate size

KEY VALUES

per 100 g as fed

Protein	40g
Fat content	10g
Crude ash	8.9g
Carbohydrate	23.9g
Dietary fibre	11.7g
Omega 6	2.47g
Omega 3	0.51g
EPA+DHA	0.28g
Calcium	1.21g
Phosphorus	1.12g
Sodium	0.9g
Linoleic acid	2.34g
L-carnitine	20mg
Metabolisable energy**	358.3kcal

COMPOSITION

Dehydrated poultry protein, maize, wheat gluten*, rice, maize gluten, hydrolysed animal proteins, vegetable fibres, animal fats, minerals, chicory pulp, fish oil, soya oil, Fructo-Oligo-Saccharides (FOS), psyllium husks and seeds, marigold extract (source of lutein).

Additives (kg) * * *

Nutritional additives: Vitamin A: 20500IU, Vitamin D3: 700IU, E1 (Iron): 42mg, E2 (Iodine): 3mg, E4 (Copper): 5mg, E5 (Manganese): 55mg, E6 (Zinc): 180mg. Preservatives - Antioxidants.

*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation

- **Calculated according to NRC2006
- ***<u>NOTE</u>: Values reflect only levels added to the formula, not those naturally occurring in components of the diet. For total levels in the diet, refer to the Average Analysis pages

FURTHER INFORMATION

Food consumption increases within 48 hours of neutering*: Changes in bodyweight can be significant, but an adapted diet may help limit weight gain.

*Nguyen PG, Dumon HJ, Siliart BS, Martin LJ, Sergheraert R, Biourge VC.

Effects of dietary fat and energy on body weight and composition after gonadectomy in cats. Am J Vet Res. 2004 Dec;65[12]:1708-13

