

FOR USE ON
SOUTHERN LAWNS
INCLUDING ALL VARIETIES OF
**ST. AUGUSTINE
GRASS**
INCLUDING
FLORATAM

BIOADVANCED
SCIENCE-BASED SOLUTIONS

SEASON LONG LAWN WEED KILLER + PREVENTER FOR SOUTHERN LAWNS

KILLS WEEDS*
*including dandelion,
clover & dollarweed*

PREVENTS Weeds*
*plus CRABGRASS
up to 6 MONTHS*

DANDELION

CLOVER

DOLLARWEED

KILLS & PREVENTS
WEEDS*
UP TO
**6
MONTHS**

*listed

TREATS UP TO
4,000 SQ. FT.
NET WEIGHT 10 lb. (4.5 kg.)

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional
precautionary statements

ACTIVE INGREDIENTS	
Dicamba acid	0.088%
Penoxsulam	0.041%
Indaziflam	0.027%
OTHER INGREDIENTS	99.844%
TOTAL	100.000%



SEASON LONG
LAWN WEED KILLER
+ PREVENTER
FOR SOUTHERN LAWNS



SEASON LONG LAWN WEED KILLER + PREVENTER

FOR SOUTHERN LAWNS

GET MORE FROM THE **BLUE BAG®**

QUICK FACTS:

Kills broadleaf weeds* including dollarweed, dandelion, clover, chickweed & henbit.

Prevents grassy and broadleaf weeds* such as crabgrass, annual bluegrass, common dandelion, goosegrass & spurge up to 6 months.

WHEN TO APPLY:
Apply anytime in early spring or fall.

WHERE TO USE:
Use only on established lawns of the turf types listed below:
Bermudagrass, Centipedegrass, all varieties of St. Augustinegrass (including Floratam) and Zoysia.

Do not use on any other turf types including: bahiagrass, bentgrass, ormond variety of bermudagrass, buffalograss, carpetgrass, dichondra, fescue, kikuyugrass, ryegrass, or seashore paspalum lawns.

COVERAGE:
This bag treats 4,000 sq. ft.

READ THE LABEL FIRST! DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

- Use this product only on established Southern lawns listed on this label – new lawns must be at least 3 months old if sodded, 16 months old if sprigged & 16 months old if seeded.
- Do not apply this product to water-saturated soils.
- Apply only to actively growing turf.
- Apply in spring after the lawn is actively growing and has been mowed 2 times.
- Do not apply while turfgrass is just turning green in early spring.
- Apply in fall no closer than 6 weeks before the first historical frost date for your area.
- Do not apply to Centipedegrass after October 1.
- Do not apply this product if the lawn appears unhealthy e.g., due to lack of water (drought, watering restrictions), turf disease (dollarspot, large patch, grey leaf spot, take-all root rot, etc.), parasitic microscopic worms (nematodes), turf insects (chinch bugs, etc.), heavy shade, recent aeration or verticutting, or if the lawn is prone to winter injury.
- Do not allow granules to come into contact with flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Apply when the air is calm to avoid drift onto desirable plants.

HOW TO USE

- Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules.
- Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back onto the lawn.
- Wait 24 hours after applying then activate residual barrier by lightly watering in.
- Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn.
- Do not make more than two applications per site in any 12 month period.
- Heavy mowing will disturb the weed preventative barrier and reduce the effectiveness of this product.

FOR BEST RESULTS

- WHEN TO APPLY:**
- Apply when - weeds are small and actively growing - temperatures are between 50-90° F.
 - May be applied to dry or (for best results) to dew-wetted lawn.
 - To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80° F range.
 - To prevent Annual Bluegrass (Poa annua) apply August - December.
 - Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

- MOWING:**
- If the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product.
 - Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

- WATERING:**
- If lawn is dry, water 1 to 2 days before using this product.
 - Do not water for at least 24 hours after application or apply if rain is expected within 24 hours.

HOW IT WORKS

This product contains a proven blend of weed killers to kill existing weeds* plus a residual herbicide to keep them from coming back for up to 6 months. This product starts working immediately to stop weed growth until weeds* are dead within weeks.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.



*listed

SPREADER SETTINGS




Fig. A – Make sure each strip overlaps the previous strip to ensure complete coverage.

Fig. B – Make sure the wheels overlap the wheel marks in the grass from the previous strip to ensure complete coverage.

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.

- Tip: 1,000 sq. ft. = 20 ft. x 50 ft.
- Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

BROADCAST SPREADER	SETTING
Earthway Ev-N-Spread	15
Sta-Green	5
Scotts 2000	4 1/2
Scotts Mini	4 3/4
Scotts Deluxe Edge Guard	4 3/4
Scotts Edge Guard Pro	4 1/4

SETTING TO DELIVER 2.5lb/1000 sq. ft.

DROP SPREADER	SETTING
Scotts Accugreen 1000	5 3/4
Scotts Accugreen 2000	6 1/2
Scotts Accugreen 3000	4 1/4

If your spreader is not listed, please call our Consumer Service Hotline toll-free 1-(877) 229-3724 for help.

KILLS			PREVENTS			KILLS AND PREVENTS				
Beggarweed, Creeping	Dollarweed (Pennywort)	Peppergrass	Ageratum, Tropic	Crabgrass, Henry	Fleabane, Hairy	Mulberry Weed	Rocket, London	Tassel Flower	Burnweed, American	Medic, Black
Beggarweed, Dwarf	Goldenrod	Pepperweed, Virginia	Barley, Mouse	Crabgrass, Large	Foxtail, Giant	Mustard, Black	Ryegrass, Italian	Velvetleaf	Chickweed, Common	Plantain, Buckhorn
Betony, Florida	Hawkweed	Plantain, Broadleaf	Barnyardgrass	(Hairy Crabgrass)	Foxtail, Green	Mustard, Wild	Ryegrass, Perennial	Vetch, Purple	Chickweed, Mouseear	Purslane, Common
Burdock	Heallal	Smartweed	Bittercress	Crabgrass, Smooth	Foxtail, Yellow	Nightshade, American	Sandbur	Wild Carrot	Clover, White	Pusley, Florida
Buttercup, Hairy	Ivy, Ground (Creeping Charlie)	Sorrel, Sheep	Black Nightshade, Hairy	Cudweed, Linear Leaf	Filaree, Redstem	Black	Sedge, Annual	Willowherb	Dandelion, Common	Ragweed, Common
Buttonweed, Virginia [†]	Knotweed	Speedwell, Common	Bluegrass, Annual	(Purple Cudweed)	Goosegrass	Nightshade, Hairy	Sesbania, Hemp	Woodsorrel	Dock, Curly	Shepardspurge
Clover, Hop	Kyllinga, Green [†]	Spotted Catsear	(Poa annua)	Dandelion, Cat's Ear	Groundsel, Common	Parthenium	Sida, Prickly (Teaweed)	(Oxalis)	Eclipta	Spurge, Spotted
Clover, Red	Lespedeza, Common	Spurweed	Brome, Foxtail	Doveweed	Guineagrass	Phyllanthus, Longstalked	Southern Brassbuttons		Henbit	Thistle, Canada
Cocklebur, Common	Mallow, Roundleaf	Thistle	Buckwheat, Wild	Evening Primrose,	Kochia	Pigweed, Prostrate	Sowthistle, Annual		Knotweed, Prostrate	Woodsorrel, Yellow
Coffeeweed,	Matchweed	Vetch, Common	Burweed, Lawn	Common	Kyllinga, Fragrant	Pigweed, Redroot	Speedwell, Corn		Kyllinga, Annual [†]	[†] Suppression
(Common Chickory)	Nutsedge, Purple [†]	Wild Violet	California Burclover	Evening Primrose,	Lovegrass, Tufted	Plantain, Paleseed	Spurge, Prostrate		Lambsquarters	
Daisy, English (Lawn Daisy)	Nutsedge, Yellow [†]	Woodsorrel, Creeping	Chamomile, False	Cutleaf	Mallow, Little	Puncturevine	Sunflower, Common			
Dandelion, False	Parsley-piert	[†] Suppression	Cheatgrass	Flatsedge, Rice	Marestail	Redmaids	Swinecress			

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:
If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions.
Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

CAUTION: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

- Harmful if swallowed or absorbed through skin.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- Remove and wash contaminated clothing before reuse.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing and wash hands.
- If product gets on skin, wash thoroughly with soap & water.
- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates, and plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.
- **Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- **Surface Water Advisory:** This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 0.088% Dicamba, (CAS # 1918-00-9), 0.041% Penoxsulam (CAS # 219714-96-2) and 0.027% Indaziflam (CAS # 950782-86-2).

ACTIVE INGREDIENTS	
Dicamba acid	0.088%
Penoxsulam	0.041%
Indaziflam	0.027%
OTHER INGREDIENTS	
TOTAL	99.844%
TOTAL	100.000%

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.



Questions or comments, call toll-free 1-(877) 229-3724 or visit us online at www.bioadvanced.com

BioAdvanced® and Get More From the Blue Bag® are registered trademarks of SBM Life Science Corp. Distributed by: SBM Life Science Corp. 1001 Winstead Drive, Suite 500, Cary, NC 27513

