

## **SEASON LONG** LAWN WEED KILLER + PREVENTER

FOR SOUTHERN LAWNS

eating, drinking, chewing gum, using tobacco, or using the toilet. • Wash thoroughly with soap and water after handling and before

• IT product gets on clothing, remove clothing as soon as possible

Flatsedge, Rice

Evening Primrose, Cutleaf

Evening Primrose,

Dandelion, Cat's Ear

• It product gets on skin, wash thoroughly with soap & water.

and launder clothing separately before reuse.

· Rinse gloves before removing and wash hands.

After using this product:

Distributed by: SBM Life Science Corp. PO Box 525, Morrisville, NC 27560 BioAdvanced® and Get More From the Blue Bag® are registered trademarks of SBM Life Science Corp. or visit us online at www.bioadvanced.com Questions or comments, call toll-free 1-(877) 229-3724

\*Or your noney back. 

performance or safety if not used according to the directions. consistent with applicable law, the Buyer assumes responsibility for lack of ions. To the extent for Use" are

	NOTICE: Research and testing have determined
100.000%	JATOT
%778.66.	OTHER INGREDIENTS
%720.0	meltizebnl
%l\$0.0	Penoxsulam
%880.0	Oicamba acid
	ACTIVE INGREDIENTS

that the "Directions	NOTICE: Research and testing have determined
100.000%	1ATOT
%448.66.	OTHER INGREDIENTS
%720.0	melfizebnl
%Iħ0.0	Penosulam malusxona9
%880.0	Dicamba acid
	ACTIVE INGREDIENTS

.(2-98-287029 # 8AO) malfizabni ntormation. Active Ingredients: 0.088% Dicamba, (CAS # 1918-00-9), 0.041% Penoxsulam (CAS # 219714-96-2) and 0.027% octor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency

nucouscions becsour Do not give anything by mouth to an

for further treatment advice. rinsing eye. the first 5 minutes, then continue

suarrenbsame

Kyllinga, Annual'

Eclipta

Dock, Curly

Slover, White

Burnweed, American

Scotts Accugreen 3000

Scotts Accugreen 2000

Scotts Accugreen 1000

SPREADER

DBUD

· Call a poison control center or doctor gently with water for 15 - 20 minutes. Hold eye open and rinse slowly and IE IN EXES

Suppression

7/L**7** 

61/5

₽/g G

SETTING

woodsorrel, Yellow

Spurge, Spotted Thistle, Canada

Pusley, Florida

Medic, Black

KILLS AND PREVENTS

SETTING TO DELIVER 2.51b/1000 sq. ft.

400217912b 180502V1

doctor for further treatment advice. of water for 15 - 20 minutes.

Vild Carrot

Vetch, Purple

43/4

43/4

7/17

G

GL

SETTING

person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

starts working immediately to stop weed growth until weeds\* are dead within weeks. residual herbicide to keep them from coming back for up to 6 months. This product

This product contains a proven blend of weed killers to kill existing weeds\* plus a

• Do not water for at least 24 hours after application or apply if rain is expected

Clippings from the first three mowings should be left on the treated area.
 Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

• It the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after

• Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

- To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80  $^\circ$  F range.

• Apply when - weeds are small and actively growing - temperatures are between 50-90° F.

FOR BEST RESULTS

GET MORE ROWTHE BLUE BAG"

• To prevent Annual Bluegrass (Poa annua) apply August - December.

 $\bullet$  May be applied to dry or (for best results) to dew-wetted lawn.

WHEN TO APPLY:

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW IT WORKS

Do not apply this product in a way that will contact any

• If lawn is dry, water 1 to 2 days before using this product.

Have person sip a glass of water if Kinse skin immediately with plenty nediately for treatment advice. Take off contaminated clothing Call a poison control center or doctor IF ON SKIN OR CLOTHING Swinecress Spurge, Prostrate Sunflower, Common Speedwell, Corn Sida, Prickly (Teaweed) (Uxalis) Sedge, Annual Gesbania, Hemp

Ryegrass, Perennial

Ryegrass, Italian

Коскет, London

spiembel Marestail Puncturevine Kyllinga, Fragrant Lovegrass, Tufted Mallow, Little Dantain, Paleseed <sup>o</sup>igweed, Redroot ssergeaning Phyllanthus, Longstalked Groundsel, Common Vightshade, American Mustard, Wild Mulberry Weed

to do so by a poison control center

Do not induce vomiting unless told

IF SWALLOWED

**GIA T2811** 

Filaree, Redstem Cudweed, Linear Leaf Hoxtail, Yellow Foxtail, Green (Hairy Crabgrass) Crabgrass, Large Crabgrass, Large Fleabane, Hairy It your spreader is not listed, please call our Consumer Service Hotline toll-free 1-(877) 229-3724 for help. Scotts Edge Guard Pro granues eveniy. the spreader works properly and spreads Scotts Deluxe Edge Guard

SPREADER SETTINGS

Scotts Mini Scotts 2000 Sta-Green Earthway Ev-N-Spread

narrow, concentrated or uneven bands. Be sure • Do not use a spreader that applies granules in • Tip: 1,000 sq. ft. = 20 ft. x 50 ft. manutacturer's directions. Calibrate your spreader according to

Settings are approximate. The condition of your

· Heavy raking will disturb the weed preventative barrier and reduce the effectiveness

• Do not overlap treated areas or make repeat applications within 90 days as over

• Wait 24 hours after applying then activate residual barrier by lightly watering in.

Sweep, brush, or blow any accidentally applied material off driveways, sidewalks

Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure
you cover lawn areas uniformly, first treat the border of the lawn then fill in the

HOW TO USE

• Do not allow granules to come into contact with flowers, ornamental shrubs or trees,

take-all root rot, etc.), parasitic microscopic worms (nematodes), turf insects (chinch bugs, etc.), heavy shade, recent aeration or verticutting, or if the lawn is

(drought, watering restrictions), turf disease (dollarspot, large patch, grey leat spot, Do not apply this product if the lawn appears unhealthy e.g., due to lack of water

Apply in fall no closer than 6 weeks before the first historical frost date for your area.

• Use this product only on established Southern lawns listed on this label – new lawns must be at least 3 months old if sodded, 16 months old if springed & 16 months old

BEFORE YOU USE

Apply in spring after the lawn is actively growing and has been mowed 2 times.

Apply when the air is calm to avoid drift onto desirable plants.

• Do not apply while turfgrass is just turning green in early spring.

ITSAIT JEBAL EHRST!

Do not apply to Centipedegrass after October 1.

Do not apply this product to water-saturated soils.

Apply only to actively growing turf.

or other desirable broadleaf plants, as serious plant injury may occur.

center. Make parallel passes, walking at a steady pace to ensure even distribution of

• Do not make more than two applications per site in any 12 month period.

previous strip to ensure complete coverage. marks in the grass from the

Cheatgrass

Chamomile, False

Burweed, Lawn California Burclover

Bluegrass, Annual (Poa

Black Nightshade,

Biftercress Barnyardgrass

Sarley, Mouse

Ageratum, Tropic

**PREVENTS** 

Buckwheat, Wild

Brome, Foxtail

strip to ensure complete artip overlaps the previous Fig. A - Make sure each

application could damage the lawn.

Fig. B – Make sure the

DROP SPREADER

**BROADCAST SPREADER** 

This bag treats 4,000 sq. ft. **COVERAGE:** 

1-0M-9778 = (37) = 071106-6A-003 (46) 1-17-7278 = (32)

This product is toxic to fish, aquatic invertebrates, and plants.

• When using this product, wear long-sleeved shirt, long pants,

PRECAUTIONARY STATEMENTS

inate water, tood, or teed by storage and disposal.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

STORAGE AND DISPOSAL

Parsley-piert

Matchweed

Mallow, Roundleaf

nsd) griqeerd) bruorê ,yvl

Dollarweed (Pennywort)

Kyllinga, Green<sup>‡</sup>

Heallal Hawkweed

ENVIRONMENTAL HAZARDS

· Kemove and wash contaminated clothing before reuse.

· Avoid contact with eyes, skin, or clothing.

storage cabinet. Protect from freezing.

Vounnun onickony) Nutsedge, Purple<sup>†</sup> Daisy, English (Lawn Daisy) Nutsedge, Yellow<sup>‡</sup> Parelou riest

Cocklebur, Common

Suttercup, Hairy

Betony, Florida

KILLS

Beggarweed, Creeping Beggarweed, Dwarf

Clover, Hop Clover, Red

Harmful if swallowed or absorbed through skin.

· causes moderate eye irritation.

surface water via runoff for several months or more after application.

EPA Reg. No. 92564-61 EPA Est. No. indicated by 3rd and 4th digits of the batch number on this package.

poorly draining soils and soils with shallow ground water. This product is classified as having a nigh potential for reaching

groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the

rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street,

waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface

CAUTION: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

PESTICIDE STORAGE: Store in original container in a cool, dry place, out of the reach of children, preferably a locked

griel, Creeping

təloiV bliW

Vetch, Common

Spotted Catsear

If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions.

Ground Water Advisory: This pesticide has properties and characteristics associated with chemicals detected in back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

• Surface Water Advisory: This pesticide may impact water quality due to runoff of rain water. This is especially true for

seashore paspalum lawns.

dichondra, fescue, kikuyugrass, ryegrass, or bermudagrass, buffalograss, carpetgrass, bahiagrass, bentgrass, ormond variety of Do not use on any other turf types including:

of St. Augustinegrass (including Floratam) Bermudagrass, Centipedegrass, all varieties types listed below: Use only on established lawns of the turf

> WHERE TO USE: Apply anytime in early spring or fall.

:YJ99A OT N3HW

dandelion, goosegrass & spurge up to 6 months. crabgrass, annual bluegrass, common

Prevents grassy and broadleaf weeds\* such as

Kills broadleaf weeds\* including dollarweed, dandelion, clover, chickweed & henbit.

**QUICK FACTS:** 

**FOR SOUTHERN LAWNS** 

+ PREVENTER

**LAWN WEED KILLER** 

SCIENCE-BASED SOLUTIONS BIOADVAUCED

**SEASON LONG** 





FOR USE ON **SOUTHERN LAWNS** INCLUDING ALL VARIETIES OF ST. AUGUSTINE

**GRASS** INCLUDING



## SEASON LONG LAWN WEED KILLER + PREVENTER

**FOR SOUTHERN LAWNS** 

DANDELION

**DOLLARWEED** 



TREATS UP TO 4,000 SQ. FT.

**NET WEIGHT 10 lb. (4.5 kg.)** 

KEEP OUT OF REACH OF CHILDREN CAUTION See back panel for additional **CLOVER**