

PAUL A. FOERSTER



Special
EDITION

Algebra AND Trigonometry

Functions and Applications

Classics Edition

Algebra AND Trigonometry

Functions and Applications

Paul A. Foerster

Paul A. Foerster has taught mathematics at Alamo Heights High School in San Antonio, Texas since 1961. In that same year he received his teaching certificate from Texas A&M University. His B.S. degree in Chemical Engineering and M.A. degree in Mathematics are from the University of Texas. Among many honors, he was awarded the Presidential Award for Excellence in Mathematics Teaching in 1983.

To A.W. Foerster, who helped me to understand the real world, to Admiral Rickover, who taught me how to write about it, and to my wife, who helps me to live in it.

©2024 by Paul A. Foerster

Veritas Press edition 2024.

Lancaster, Pennsylvania

www.veritaspress.com

First published by Addison-Wesley Publishing Company.

ISBN 978-1-956402-41-4

All rights reserved. No part of this book may be reproduced without permission from Veritas Press, except by a reviewer who may quote brief passages in a review; nor may any part of this book be reproduced, stored in a retrieval system or transmitted in any form by any means, electronic, mechanical, photocopying, recording or otherwise, without prior, written permission from Veritas Press.


Printed in the United States of America.

Foreword

Algebra and Trigonometry—Functions and Applications is designed for a course in intermediate algebra, advanced algebra, and trigonometry. The book can be used in two different ways:

1. As an algebra and trigonometry book, with applications,
2. As an applications book, with supporting algebra and trigonometry.

In either case, there are two possible sequences of presentation:



Applications are handled by creating mathematical models of phenomena in the real world. Students must select a kind of function that fits a given situation, and derive an equation that suits the information in the problem. The equation is then used to predict values of y when x is given or values of x when y is given. Sometimes students must use the results of their work to make interpretations about the real world, such as what “slope” means, or why there cannot be people as small as in *Gulliver’s Travels*. The problems require the students to use *many* mathematical concepts in the *same* problem. This is in contrast to the traditional “word problems”

of elementary algebra, in which the same one concept is used in many problems.

The computer is assumed to be a normal part of students' classroom experience rather than simply a novelty. For the most part, students are expected to use existing computer programs, notably computer graphic programs, rather than write their own. A disk accompanying the Teacher's Resource Book contains programs written by the author that are sufficient for the purposes of this text. You are, however, urged to seek out commercially-available software that is faster and more user-friendly.

Ideas for using calculators have caused substantial differences in presentation of certain topics. For instance, exponential equations are solved at the beginning of Chapter 6 by iterative methods on the calculators. Logarithms then arise naturally as a quicker way to get the unknown exponent. Presented this way, there is no doubt in the students' minds that a logarithm is an exponent! The calculator thus leads to a better understanding of theoretical concepts and is not simply a way to work old problems quicker.

Pedagogically, there are many opportunities for review of previous concepts. One major way this review is accomplished is through the "Do These Quickly" problems at the beginning of each problem set. Once the students have learned to work a particular kind of problem, they develop speed as these problems reappear in five-minute exercises that concentrate only on answers. Mathematical-model problems have been spread out so the students must recall how to use linear and quadratic functions while they are working exponential model problems.

Work on data analysis is included in some problem sets to show students how to decide which kind of function is an appropriate model. Material on statistics, including the normal distribution, appears in the probability chapter.

The text still starts with a brief review of the basic axioms and properties. It moves quickly to topics the students have probably not seen before, at least in the method of presentation. The purpose is two-fold. First, students should not feel as if most of the course is spent reviewing elementary algebra. Second, the presentation emphasizes the role of algebra and trigonometry as the foundation for calculus, rather than as the completion of elementary algebra. By presenting both algebra and trigonometry as the study of classes of functions, students learn the essential unity of the two subjects.

Henry Pollack of Bell Telephone Labs claims that there are just two kinds of numbers: “real” numbers such as encountered in everyday life, and “fake” numbers such as encountered in most mathematics classes! Since this book has many problems involving untidy decimals (“real” numbers), a calculator or computer are called for where appropriate. There are also problems that have small-integer answers (“fake” numbers) so that students may gain confidence in their work when they are just learning a new technique.

Since all educators share the responsibility of teaching students to read and write, there are discovery exercises so that students may wrestle with a new concept before it is reinforced by classroom discussion. The students are helped with this reading by the fact that much of the wording came from the mouths of my own students. Special thanks go to Susan Cook, Brad Foster, and Nancy Carnes, whose good class notes supplied input for certain sections. Students Lewis Donzis and David Fey wrote computer programs for some of the problems.

Thanks go to instructors in Florida, Illinois, Pennsylvania, South Dakota, Texas, and Virginia for pilot testing the original materials. Special thanks go to Bob Enenstein and his instructors in California for pilot testing the second edition. The text reflects comments from review and classroom testing by Charley Brown, Sharon Sasch Button, Pat Causey, Loyce Collenback, Bob Davies, Walter DeBill, Rich Dubsky, Michelle Edge, Sandra Frasier, Byron Gill, Pat Johnson, Michael Keeton, Carol Kipps, Bill McNabb, Shirley Scheiner, Ann Singleton, Chuck Straley, Rhett Tatsch, Joel Teller, Susan Thomas, Kay Thompson, Zalman Usiskin, Jim Wieboldt, Marv Wielard, Mercille Wisakowsky, Martha Zelinka, and Isabel Zsohar. Calvin Butterball and Phoebe Small appear with the kind permission of their parents, Richard and Josephine Andree.

Paul A. Foerster

Contents

1 PRELIMINARY INFORMATION 1

- 1-1 Sets of Numbers 1
- 1-2 The Field Axioms 3
- 1-3 Variables and Expressions 9
- 1-4 Polynomials 16
- 1-5 Equations 20
- 1-6 Inequalities 27
- 1-7 Properties Provable from the Axioms 32
- 1-8 Chapter Review and Test 44



number line

2 FUNCTIONS AND RELATIONS 50

- 2-1 Graphs of Equations with Two Variables 51
- 2-2 Graphs of Functions 53
- 2-3 Functions in the Real World 58
- 2-4 Graphs of Functions and Relations 64
- 2-5 Chapter Review and Test 69



dependent and
independent variables

3 LINEAR FUNCTIONS 72

- 3-1 Introduction to Linear Functions 73
- 3-2 Properties of Linear Function Graphs 74
- 3-3 Other Forms of the Linear Function Equation 82
- 3-4 Equations of Linear Functions from Their Graphs 86
- 3-5 Linear Functions as Mathematical Models 93
- 3-6 Chapter Review and Test 105



linear functions with
positive and negative
slopes

4 SYSTEMS OF LINEAR EQUATIONS AND INEQUALITIES 110

- 4-1 Introduction to Linear Systems 111
- 4-2 Solution of Systems in Linear Equations 112
- 4-3 Second-Order Determinants 120
- 4-4 $f(x)$ Terminology, and Systems as Models 126
- 4-5 Linear Equations with Three or More Variables 133
- 4-6 Systems of Linear Equations with Three or More Variables 138
- 4-7 Solution of Second-Order Systems by Augmented Matrices 142
- 4-8 Solution of Higher-Order Systems by Augmented Matrices 146



intersection of linear
functions

- 4-9 Higher-Order Determinants 150
- 4-10 Systems of Linear Inequalities 153
- 4-11 Linear Programming 157
- 4-12 Chapter Review and Test 169



parabolas with positive
and negative x^2
coefficient

5 QUADRATIC FUNCTIONS AND COMPLEX NUMBERS 173

- 5-1 Introduction to Quadratic Functions 174
- 5-2 Graphs of Quadratic Functions 175
- 5-3 x -Intercepts and the Quadratic Formula 181
- 5-4 Imaginary and Complex Numbers 188
- 5-5 Evaluating Quadratic Functions 195
- 5-6 Equations of Quadratic Functions from Their Graphs 200
- 5-7 Quadratic and Linear Functions as Mathematical Models 204
- 5-8 Chapter Review and Test 221
- 5-9 Cumulative Review: Chapters 1 through 5 224



increasing and
decreasing exponential
functions

6 EXPONENTIAL AND LOGARITHMIC FUNCTIONS 228

- 6-1 Introduction to Exponential Functions 229
- 6-2 Exponentiation for Positive Integer Exponents 230
- 6-3 Properties of Exponentiation 236
- 6-4 Exponentiation for Rational Exponents 241
- 6-5 Powers and Radicals Without Calculators 248
- 6-6 Scientific Notation 253
- 6-7 Exponential Equations Solved by Brute Force 260
- 6-8 Exponential Equations Solved by Logarithms 263
- 6-9 Logarithms with Other Bases 269
- 6-10 Properties of Logarithms 274
- 6-11 Proofs of Properties of Logarithms 282
- 6-12 Inverses of Functions—The Logarithmic Function 289
- 6-13 The Add-Multiply Property of Exponential Functions 295
- 6-14 Exponential and Other Functions as Mathematical Models 300
- 6-15 Chapter Review and Test 317



hyperbola with
removable discontinuity

7 RATIONAL ALGEBRAIC FUNCTIONS 322

- 7-1 Introduction to Rational Algebraic Functions 323
- 7-2 Rational Function Graphs—Discontinuities and Asymptotes 324
- 7-3 Special Products and Factoring 328
- 7-4 More Factoring and Graphing 336
- 7-5 Long Division of Polynomials 343

- 7-6 Factoring Higher-Degree Polynomials—The Factor Theorem 347
- 7-7 Products and Quotients of Rational Expressions 356
- 7-8 Sums and Differences of Rational Expressions 364
- 7-9 Graphs of Rational Algebraic Functions, Again 370
- 7-10 Fractional Equations and Extraneous Solutions 378
- 7-11 Variation Functions 383
- 7-12 Chapter Review and Test 405

8 IRRATIONAL ALGEBRAIC FUNCTIONS 412

- 8-1 Introduction to Irrational Algebraic Functions 413
- 8-2 Graphs of Irrational Functions 414
- 8-3 Radicals and Simple Radical Form 416
- 8-4 Radical Equations 425
- 8-5 Variation Functions with Non-Integer Exponents 432
- 8-6 Functions of More Than One Independent Variable 442
- 8-7 Chapter Review and Test 452
- 8-8 Cumulative Review: Chapters 6 through 8 455



direct power variation

9 QUADRATIC RELATIONS AND SYSTEMS 460

- 9-1 Introduction to Quadratic Relations 461
- 9-2 Circles 462
- 9-3 Ellipses 468
- 9-4 Hyperbolas 477
- 9-5 Parabolas 486
- 9-6 Equations from Geometrical Definitions 490
- 9-7 Quadratic Relations— xy -Term 495
- 9-8 Systems of Quadratics 497
- 9-9 Chapter Review and Test 507



ellipse and hyperbola

10 HIGHER-DEGREE FUNCTIONS AND COMPLEX NUMBERS 513

- 10-1 Introduction to Higher-Degree Functions 514
- 10-2 Complex Number Review 515
- 10-3 Quadratic Equations from Their Solutions—Complex Number Factors 523
- 10-4 Graphs of Higher-Degree Functions—Synthetic Substitution 530
- 10-5 Descartes' Rule of Signs and the Upper Bound Theorem 544
- 10-6 Higher-Degree Functions as Mathematical Models 549
- 10-7 Chapter Review and Test 554



cubic function



linear and geometric series

11 SEQUENCES AND SERIES 559

- 11-1 Introduction to Sequences 560
- 11-2 Arithmetic and Geometric Sequences 564
- 11-3 Arithmetic and Geometric Means 572
- 11-4 Introduction to Series 577
- 11-5 Arithmetic and Geometric Series 581
- 11-6 Convergent Geometric Series 589
- 11-7 Sequences and Series as Mathematical Models 597
- 11-8 Factorials 612
- 11-9 Introduction to Binomial Series 617
- 11-10 The Binomial Formula 619
- 11-11 Chapter Review and Test 625



bell-shaped curve

12 PROBABILITY, DATA ANALYSIS, AND FUNCTIONS OF A RANDOM VARIABLE 631

- 12-1 Introduction to Probability 632
- 12-2 Words Associated with Probability 633
- 12-3 Two Counting Principles 636
- 12-4 Probabilities of Various Permutations 642
- 12-5 Probabilities of Various Combinations 651
- 12-6 Properties of Probability 661
- 12-7 Functions of a Random Variable 671
- 12-8 Mathematical Expectation 681
- 12-9 Statistics and Data Analysis 690
- 12-10 Chapter Review and Test 700
- 12-11 Cumulative Review: Chapters 9 through 12 705



cosine, tangent, and secant

13 TRIGONOMETRIC AND CIRCULAR FUNCTIONS 709

- 13-1 Introduction to Periodic Functions 710
- 13-2 Measurement of Arcs and Rotation 711
- 13-3 Definitions of Trigonometric and Circular Functions 717
- 13-4 Approximate Values of Trigonometric and Circular Functions 729
- 13-5 Graphs of Trigonometric and Circular Functions 735
- 13-6 General Sinusoidal Graphs 742
- 13-7 Equations of Sinusoids from Their Graphs 750
- 13-8 Sinusoidal Functions as Mathematical Models 756
- 13-9 Inverse Circular Functions 767
- 13-10 Evaluation of Inverse Relations 781
- 13-11 Inverse Circular Relations as Mathematical Models 786
- 13-12 Chapter Review and Test 793

14 PROPERTIES OF TRIGONOMETRIC AND CIRCULAR FUNCTIONS 799

- 14-1 Three Properties of Trigonometric Functions 800
- 14-2 Trigonometric Identities 807
- 14-3 Properties Involving Functions of More Than One Argument 814
- 14-4 Multiple-Argument Properties 825
- 14-5 Half-Argument Properties 831
- 14-6 Sum and Product Properties 837
- 14-7 Linear Combination of Cosine and Sine with Equal Arguments 841
- 14-8 Simplification of Trigonometric Expressions 845
- 14-9 Trigonometric Equations 850
- 14-10 Chapter Review and Test 858



secant, cosine, and
tangent

15 TRIANGLE PROBLEMS 862

- 15-1 Right Triangle Problems 863
- 15-2 Oblique Triangles—Law of Cosines 873
- 15-3 Area of a Triangle 880
- 15-4 Oblique Triangles—Law of Sines 883
- 15-5 The Ambiguous Case 888
- 15-6 General Solution of Triangles 894
- 15-7 Vectors 898
- 15-8 Vectors—Resolution into Components 906
- 15-9 Real-World Triangle Problems 915
- 15-10 Chapter Review and Test 923
- 15-11 Cumulative Review: Chapters 13 through 15 927



vector addition

FINAL EXAMINATION 930**APPENDIX A OPERATIONS WITH MATRICES 937****APPENDIX B MATHEMATICAL INDUCTION 944****APPENDIX C AN EXPLORATION OF FUNCTIONS 957****SKILLS HANDBOOK 1047**

A resource for problem solving strategies and other skills you may want to review during this course.

TABLES 965

I	Squares and Square Roots, Cubes and Cube Roots	965
II	Four-Place Logarithms of Numbers	967
III	Trigonometric Functions and Degrees-to-Radians	969
IV	Circular Functions and Radians-to-Degrees	976


GLOSSARY 979

ANSWERS TO SELECTED PROBLEMS 983

PROBLEM TITLE INDEX 1031

GENERAL INDEX 1035

1

Preliminary
Information

*Contrary to former views of mathematics, numbers were **invented** by people, rather than simply being discovered. In this book you will see how things invented mainly to form a complete mathematical system can be used to describe things that happen in the real world. First, however, you must be sure that you and your instructor are speaking the same language! The first chapter is designed with this purpose in mind.*

From previous work in mathematics you should recall the names of different kinds of numbers (positive, even, irrational, etc.). In this section you will refresh your memory so that you will know the exact meaning of these names.

Objective:

Given the name of a set of numbers, provide an example; or given a number, name the sets to which it belongs.

There are two major sets of numbers you will deal with in this course, the *real* numbers and the *imaginary* numbers. The real numbers are given this name because they are used for “real” things such as measuring and counting. The imaginary numbers are square roots of negative numbers. They are useful, too, but you must learn more mathematics to see why.

The real numbers are all numbers which you can plot on a number line (see Figure 1-1). They can be broken into subsets in several ways. For instance, there are positive and negative real numbers, integers and non-integers, rational and irrational real numbers, and so forth. The diagram facing this page shows some subsets of the set of real numbers.

The numbers in the diagram were invented in *reverse* order. The natural (or “counting”) numbers came first because mathematics was first used for counting. The negative numbers (those less than zero) were invented so that there would always be answers to subtraction problems. The rational numbers were invented to provide answers to division problems, and the irrational ones came when it was shown that numbers such as $\sqrt{2}$ could not be expressed as a ratio of two integers.



The real number line



Other operations you will invent, such as taking logarithms and cosines, lead to irrational numbers which go beyond even extracting roots. These are called “transcendental” numbers, meaning “going beyond.” When all of these various kinds of numbers are put together, you get the set of *real* numbers. The imaginary numbers were invented because no real number squared equals a negative number. Later, you will see that the real and imaginary numbers are themselves simply subsets of a larger set, called the “complex numbers.”

The following exercise is designed to help you accomplish the objectives of this section.

EXERCISE 1-1

- Write a definition for each of the following sets of numbers. Try to do this *without* referring to the diagram opposite page 1. Then look to make sure you are correct.
 - {integers}
 - {even numbers}
 - {negative numbers}
 - {irrational numbers}
 - {real numbers}
 - {counting numbers}
 - {digits}
 - {positive numbers}
 - {rational numbers}
 - {imaginary numbers}
 - {natural numbers}
 - {transcendental nos.}
- Write an example of each type of number mentioned in Problem 1.
- Copy the chart at right. Put a check mark in each box for which the number on the left of the chart belongs to the set across the top.
- Write another name for {natural numbers}.
- Which of the sets of numbers in Problem 1 do you suppose was the *first* to be invented? Why?
- One of the sets of numbers in Problem 1 contains all but one of the others as subsets.
 - Which one *contains* the others?
 - Which one is left out?
- Do decimals such as 2.718 represent *rational* numbers or *irrational* numbers? Explain.
- Do repeating decimals such as 2.3333 . . . represent *rational* numbers or *irrational* numbers? Explain.
- What real number is neither positive nor negative?

	Integers	Digits	Even Numbers	Positive Numbers	Negative Numbers	Rational Numbers	Irrational Numbers	Imaginary Numbers	Real Numbers	Natural Numbers	Counting Numbers	Transcendental Numbers
a. 5												
b. $2/3$												
c. -7												
d. $\sqrt{3}$												
e. $\sqrt{16}$												
f. $\sqrt{-16}$												
g. $\sqrt{-15}$												
h. 44												
i. π												
j. 1.765												
k. -10000												
l. $-1\frac{1}{2}$												
m. $-\sqrt{6}$												
n. 0												
o. 1												
p. $1/9$												

1-2 THE FIELD AXIOMS

From previous mathematics courses you probably remember names such as “Distributive Property,” “Reflexive Property,” and “Multiplication Property of Zero.” Some of these properties, called *axioms*, are accepted without proof and are used as starting points for working with numbers.



From a small number of rather obvious axioms, you will derive all the other properties you will need. In this section you will concentrate on the axioms that apply to the *operations* with numbers such as $+$ and $\times$. In Section 1-7 you will find the axioms that apply to the *relationships* between numbers, such as $=$ and $<$.

Objective:

Given the name of an axiom that applies to $+$ or $\times$, give an example that shows you understand the meaning of the axiom; and vice versa.

There are eleven axioms that apply to adding and multiplying real numbers. These are called the *Field Axioms*, and are listed in the following table. If you already feel familiar with these axioms, you may go right to the problems in Exercise 1-2. If not, then read on!

THE FIELD AXIOMS

CLOSURE

$\{\text{real numbers}\}$ is *closed* under addition and under multiplication. That is, if x and y are real numbers, then

$x + y$ is a *unique, real* number,
 xy is a *unique, real* number.

COMMUTATIVITY

Addition and multiplication of real numbers are *commutative* operations. That is, if x and y are real numbers, then

$x + y$ and $y + x$ are *equal* to each other,
 xy and yx are equal to each other.

ASSOCIATIVITY

Addition and multiplication of real numbers are *associative* operations. That is, if x , y , and z are real numbers, then

$(x + y) + z$ and $x + (y + z)$ are *equal* to each other.
 $(xy)z$ and $x(yz)$ are *equal* to each other.

DISTRIBUTIVITY

Multiplication *distributes* over addition. That is, if x , y , and z are real numbers, then

$x(y + z)$ and $xy + xz$ are *equal* to each other.

IDENTITY ELEMENTS

$\{\text{real numbers}\}$ contains:

A *unique* identity element for *addition*, namely 0. (Because $x + 0 = x$ for any real number x .)

A *unique* identity element for *multiplication*, namely 1. (Because $x \cdot 1 = x$ for any real number x .)

INVERSES

{real numbers} contains:

A *unique additive* inverse for every real number x . (Meaning that every real number x has a real number $-x$ such that $x + (-x) = 0$.)

A *unique multiplicative* inverse for every real number x except zero.

(Meaning that every non-zero number x has a real number $\frac{1}{x}$ such that $x \cdot \frac{1}{x} = 1$.)

Notes:

1. Any set that obeys all eleven of these axioms is a *field*.
2. The eleven Field Axioms come in 5 pairs, one of each pair being for addition and the other for multiplication. The Distributive Axiom expresses a relationship between these two operations.
3. The properties $x + 0 = x$ and $x \cdot 1 = x$ are sometimes called the "Addition Property of 0" and the "Multiplication Property of 1," respectively, for obvious reasons.
4. The number $-x$ is called, "the *opposite* of x ," "the *additive inverse* of x ," or "negative x ."
5. The number $\frac{1}{x}$ is called the "multiplicative inverse of x ," or the "reciprocal of x ."

Closure—By saying that a set is "closed" under an operation, you mean that you cannot get an answer that is *out* of the set by performing that operation on numbers *in* the set. For example, $\{0, 1\}$ is closed under multiplication because $0 \times 0 = 0$, $0 \times 1 = 0$, $1 \times 0 = 0$, and $1 \times 1 = 1$. All the answers are *unique*, and are *in* the given set. This set is *not* closed under addition because $1 + 1 = 2$, and 2 is *not* in the set. It is not closed under the operation "taking the square root" since there are *two* different square roots of 1: $+1$ and -1 .

Commutativity—The word "commute" comes from the Latin word "commutare," which means "to exchange." People who travel back and forth between home and work are called "commuters" because they regularly exchange positions. The fact that addition and multiplication are commutative operations is somewhat unusual. Many operations such as subtraction and exponentiation (raising to powers) are *not* commutative. For example,

$$2 - 5 \text{ does not equal } 5 - 2,$$

and

$$2^3 \text{ does not equal } 3^2.$$



Indeed, most operations in the real world are not commutative. Putting on your shoes and socks (in that order) produces a far different result from putting on your socks and shoes!

Associativity—You can remember what this axiom states by remembering that to “associate” means to “group.” Addition and multiplication are associative, as shown by

$$(2 + 3) + 4 = 9 \quad \text{and} \quad 2 + (3 + 4) = 9.$$

But subtraction is *not* associative. For example,

$$(2 - 3) - 4 = -5 \quad \text{and} \quad 2 - (3 - 4) = 3.$$

Distributivity—Parentheses in an expression such as $2 \times (3 + 4)$ mean, “Do what is inside *first*.” But you don’t *have* to do $3 + 4$ first. You could “distribute” a 2 to each term inside the parentheses, getting $2 \times 3 + 2 \times 4$. The Distributive Axiom expresses the fact that you get the same answer either way. That is,

$$2 \times (3 + 4) = 14 \quad \text{and} \quad 2 \times 3 + 2 \times 4 = 14.$$

Note that multiplication does *not* distribute over multiplication. For example,

$$2 \times (3 \times 4) \quad \text{does not equal} \quad 2 \times 3 \times 2 \times 4,$$

as you can easily check by doing the arithmetic.

Identity Elements—The numbers 0 and 1 are called “identity elements” for adding and multiplying, respectively, since a number comes out “identical” if you add 0 or multiply by 1. For example,

$$5 + 0 = 5 \quad \text{and} \quad 5 \times 1 = 5.$$

Inverses—A number is said to be an *inverse* of another number for a certain operation if it “undoes” (or inverts) what the other number did. For example, $\frac{1}{3}$ is the multiplicative inverse of 3. If you start with 5 and multiply by 3 you get

$$5 \times 3 = 15.$$

Multiplying the answer, 15, by $\frac{1}{3}$ gives

$$15 \times \frac{1}{3} = 5,$$

which “undoes” or “inverts” the multiplication by 3. It is easy to tell if two numbers are *multiplicative inverses* of each other because their product is always equal to 1, the multiplicative identity element. For example,

$$3 \times \frac{1}{3} = 1.$$

Similarly, two numbers are *additive inverses* of each other if adding them to each other gives 0, the additive identity element. For example, $\frac{5}{7}$ and $-\frac{5}{7}$ are additive inverses of each other because

$$\frac{5}{7} + (-\frac{5}{7}) = 0.$$

The following exercise is designed to familiarize you with the names and meanings of the Field Axioms.

EXERCISE 1-2

Do These Quickly

The following problems are intended to refresh your skills. Some problems come from the last section, and others probe your general knowledge of mathematics. You should be able to do all 10 in less than 5 minutes.

- Q1. Simplify: $11 - 3 + 5$
- Q2. Multiply and simplify: $\left(\frac{2}{3}\right)\left(\frac{6}{7}\right)$
- Q3. Add: $3.74 + 5$
- Q4. If $x + 7$ is 42, what does x equal?
- Q5. Is -13 an integer?
- Q6. Multiply: $(9x)(6x)$
- Q7. Square 7.
- Q8. Is 1.3 a rational number?
- Q9. Multiply: $5(3x - 8)$
- Q10. Simplify: $(-3)(0.7)(-5)(-1)$

Work the following problems.

1. Tell what is meant by
 - a. additive identity element,
 - b. multiplicative identity element.
2. What is
 - a. the *additive* inverse of $\frac{2}{3}$?
 - b. the *multiplicative* inverse of $\frac{2}{3}$?
3. Using variables (x , y , z , etc.) to stand for numbers, write an example of each of the eleven field axioms. Try to do this by writing all eleven



examples first, then checking to be sure you are right. Correct any which you left out or got wrong.

4. Explain why 0 has *no* multiplicative inverse.
5. The Closure Axiom states that you get a *unique* answer when you add two real numbers. What is meant by a “unique” answer?
6. You get the same answer when you add a column of numbers “up” as you do when you add it “down.” What axiom(s) show that this is true?
7. Calvin Butterball and Phoebe Small use the distributive property as follows:

$$\text{Calvin: } 3(x + 4)(x + 7) = (3x + 12)(x + 7).$$

$$\text{Phoebe: } 3(x + 4)(x + 7) = (3x + 12)(3x + 21).$$

Who is right? What mistake did the other one make?

8. Write an example which shows that:
 - a. Subtraction is *not* a commutative operation.
 - b. {negative numbers} is *not* closed under multiplication.
 - c. {digits} is *not* closed under addition.
 - d. {real numbers} is *not* closed under the $\sqrt{\quad}$ operation (taking the square root).
 - e. Exponentiation (“raising to powers”) is *not* an associative operation. (Try 4^{2^3} .)
9. For each of the following, tell which of the Field Axioms was used, and whether it was an axiom for *addition* or for *multiplication*. Assume that x , y , and z stand for real numbers.
 - a. $x + (y + z) = (x + y) + z$
 - b. $x \cdot (y + z)$ is a real number
 - c. $x \cdot (y + z) = x \cdot (z + y)$
 - d. $x \cdot (y + z) = (y + z) \cdot x$
 - e. $x \cdot (y + z) = xy + xz$
 - f. $x \cdot (y + z) = x \cdot (y + z) + 0$
 - g. $x \cdot (y + z) + (-[x \cdot (y + z)]) = 0$
 - h. $x \cdot (y + z) = x \cdot (y + z) \cdot 1$
 - i. $x \cdot (y + z) \cdot \frac{1}{x \cdot (y + z)} = 1$
10. Tell whether or not the following sets are *fields* under the operations $+$ and $\times$. If the set is not a field, tell which one(s) of the Field Axioms do not apply.
 - a. {rational numbers}
 - b. {integers}
 - c. {positive numbers}
 - d. {non-negative numbers}

1-3 | VARIABLES AND EXPRESSIONS

In previous mathematics courses you have seen *expressions*, such as

$$3x^2 + 5x - 7,$$

that stand for numbers. Just *what* number an expression stands for depends on what value you pick for the *variable* (x in this case). The name “variable” is picked because x can stand for various different numbers at different times. The numbers 3, 5, -7 , $\frac{9}{10}$, $\sqrt{11}$ etc., are called *constants* because they stand for the *same* number *all* the time.

In this section you will *evaluate* expressions by substituting values for the variable. In order to do this more easily, you can *simplify* the expression using the axioms of the previous section.

Objective:

Given an expression containing a variable,

- evaluate* it by substituting a given number for the variable, and finding the value of the expression,
- simplify* it by using the Field Axioms to transform it to an equivalent expression that is easier to evaluate.

DEFINITION**VARIABLE**

A **variable** is a letter which stands for an *unspecified* number from a *given* set.

For example, if the set you have in mind is {digits}, and x is the variable, then x could stand for any one of the numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, or 9. In this case, {digits} is called the *domain* of x . The word comes from the Latin “domus,” meaning “house.” So the domain of a variable is “where it lives.” Since the domain of most variables in this course will be {real numbers}, you make the following agreement:

AGREEMENT

Unless otherwise specified, the domain of a variable will be assumed to be the set of all real numbers.



DEFINITION

EXPRESSION

An **expression** is a collection of variables and constants connected by operation signs (+, −, ×, ÷, etc.) which stands for a *number*.

To find out *what* number an expression stands for, you must substitute a value for each variable, then do the indicated operations.

EXAMPLE 1

Evaluate $3x^2 + 5x - 7$ if $x = 4$.

Solution:

$$\begin{aligned}
 & 3x^2 + 5x - 7 && \text{Write the given expression.} \\
 = & 3 \cdot 4^2 + 5 \cdot 4 - 7 && \text{Substitute 4 for } x. \\
 = & 3 \cdot 16 + 5 \cdot 4 - 7 && \text{Square the 4.} \\
 = & 48 + 20 - 7 && \text{Do the multiplication.} \\
 = & \underline{61} && \text{Add and subtract from left to right.}
 \end{aligned}$$

There are several things you should realize about the preceding calculations. First, you must substitute the *same* value of x *everywhere* it appears in the expression. Although a variable can take on different values at different times, it stands for the *same* number at any *one* time. This fact is expressed in the Reflexive Axiom, which states, “ $x = x$.”

The second thing you should realize is that this expression involves *subtraction* and *exponentiation* (raising to powers). These operations, as well as *division*, can be defined in terms of addition and multiplication.

DEFINITIONS

Subtraction: $x - y$ means $x + (-y)$.

Division: $x \div y$ means $x \cdot \frac{1}{y}$. (The symbols $\frac{x}{y}$ and x/y are also used for $x \div y$.)

Exponentiation: x^n means n , x 's *multiplied* together. For example,

$$x^3 \text{ means } x \cdot x \cdot x.$$

The third thing you should realize is that the answer you get depends on the *order* in which you do the operations. So that there will be no doubt about what an expression such as $3x^2 + 5x - 7$ means, you make the following agreement:

AGREEMENT

ORDER OF OPERATIONS

1. Do any operations inside parentheses *first*.
2. Do any exponentiating next.
3. Do multiplication and division in the order in which they occur, from left to right.
4. Do addition and subtraction last, in the order in which they occur, from left to right.

EXAMPLE 2

Carry out the following operations:

- a. $3 + 4 \times 5$ Multiply *first*.
 $= 3 + 20$
 $= \underline{\underline{23}}$ Add *last*.
- b. $3 + 4 \times 5 \div 2$
 $= 3 + 20 \div 2$ Multiply and divide from left to right.
 $= 3 + 10$ Divide *before* adding.
 $= \underline{\underline{13}}$ Add *last*.
- c. $3 - 4 \times 5 \div 2 + 9$
 $= 3 - 20 \div 2 + 9$ Multiply and divide from left to right.
 $= 3 - 10 + 9$ Divide *before* $+$ and $-$.
 $= -7 + 9$ Add and subtract from left to right.
 $= \underline{\underline{2}}$ Add and subtract last. ■

An expression might contain the *absolute value* operation. The symbol $|x|$ means the *distance* between the number x and the origin of the number line. For example, $|-3|$ and $|3|$ are both equal to 3, since both 3 and -3 are located 3 units from the origin (Figure 1-3).




Figure 1-3

Similarly,

$$|5| = 5$$

$$|-7| = 7$$

$$|0| = 0,$$

and so forth.

The absolute value of a *variable* presents a problem. If x is a *positive* number, then $|x|$ is equal to x . But if x is a *negative* number, then $|x|$ is equal to the *opposite* of x . For instance, if $x = -9$, then

$$|x| = |-9| = -(-9) = 9.$$

A precise definition of absolute value can be written as follows:

DEFINITION

$$|x| = x \text{ if } x \text{ is positive (or 0)}$$

$$|x| = -x \text{ if } x \text{ is negative}$$

EXAMPLE 3

Evaluate $|17 - 4x| - 2$ if

- $x = 5$
- $x = -3$.

- $$\begin{aligned}
 &|17 - 4x| - 2 \\
 &= |17 - 20| - 2 && \text{Substitute 5 for } x. \\
 &= |-3| - 2 && \text{Arithmetic} \\
 &= 3 - 2 && \text{Definition of absolute value} \\
 &= \underline{1} && \text{Arithmetic}
 \end{aligned}$$
- $$\begin{aligned}
 &|17 - 4x| - 2 \\
 &= |17 + 12| - 2 && \text{Substitute } -3 \text{ for } x.
 \end{aligned}$$

$$\begin{aligned}
 &= |29| - 2 && \text{Arithmetic} \\
 &= 29 - 2 && \text{Definition of absolute value} \\
 &= \underline{\underline{27}} && \text{Arithmetic}
 \end{aligned}$$

Two expressions are equivalent if they *equal* each other for *all* values of the variable. For example, $3x + 8x$ and $11x$ are equivalent expressions. *Simplifying* an expression means transforming it to an equivalent expression that is in some way simpler to work with. The expression $11x$ is considered to be simpler than $3x + 8x$ because it is easier to evaluate when you pick a value of x . Adding the $3x$ and $8x$ is called “collecting like terms.” It is justified by using the Distributive Axiom *backwards*.

$$\begin{aligned}
 3x + 8x &= (3 + 8)x && \text{Distributivity} \\
 &= 11x && \text{Arithmetic}
 \end{aligned}$$

EXAMPLE 4

Simplify $7x \cdot 2 \div x$.

Since the Field Axioms apply to multiplication rather than division, you would treat “ $\div x$ ” as “ $\cdot \frac{1}{x}$ ”, commute the multiplication, and get

$$\begin{aligned}
 &7x \cdot 2 \div x \\
 &= 7x \cdot 2 \cdot \frac{1}{x} && \text{Definition of division} \\
 &= \left(7x \cdot \frac{1}{x}\right) \cdot 2 && \text{Commutativity and associativity} \\
 &= 7 \cdot 2 && \text{Associativity and multiplicative inverses} \\
 &= \underline{\underline{14}} && \text{Arithmetic}
 \end{aligned}$$

EXAMPLE 5

Simplify $2 - 3[x - 2 - 5(x - 1)]$.

Here you must observe the agreed-upon sequence of operations. The first thing to do is *start inside the innermost parentheses* and work your way out (like a termite!).

$$\begin{aligned}
 &2 - 3[x - 2 - 5(x - 1)] \\
 &= 2 - 3[x - 2 - 5x + 5] && \text{Distributivity} \\
 &= 2 - 3[-4x + 3] && \text{Collecting like terms} \\
 &= 2 + 12x - 9 && \text{Distributivity} \\
 &= \underline{\underline{12x - 7}} && \text{Commutativity and associativity}
 \end{aligned}$$



Notes:

1. You must remember some things from previous mathematics courses. For example, a negative number times a negative number is a *positive* number. This sort of thing can be proved using the Field Axioms, as you will see in Section 1-7.
2. There are several kinds of symbols of inclusion.
 - () Parentheses.
 - [] Brackets.
 - { } Braces (also used for *set* symbols).
 - Vinculum (an overhead line, used in fractions and elsewhere, such as in $\frac{x-3}{x+7}$).

To avoid so many different symbols, sometimes “nested” parentheses are used. For example, the expression

$$2 - (3 + 4(5 - 6(7 + x)))$$

would be simplified by starting with the *innermost* parentheses.

In the following exercise, you will practice simplifying and evaluating expressions. If the going gets difficult, just tell yourself that no matter how complicated an expression looks, it just stands for a *number*. And people *invented* numbers!

EXERCISE 1-3

Do These Quickly

The following problems are intended to refresh your skills. Some are from the first two sections of this chapter, and others probe your general knowledge of mathematics. You should be able to do all 10 in less than 5 minutes.

- Q1. Is $\sqrt{9}$ an integer?
- Q2. Is $-\frac{4}{7}$ a real number?
- Q3. Commute the 3 and the x : $2y + 3 + x$
- Q4. Associate the $4a$ and the $2c$: $4a + 2c + 5d$
- Q5. Distribute the 5: $5(3x - 7)$
- Q6. Write the additive inverse of $\frac{5}{8}$.

- Q7. Write the multiplicative identity element.
- Q8. If $3x$ equals 42, what does x equal?
- Q9. Multiply: $(2.3)(4)$
- Q10. Divide and simplify: $(\frac{2}{3}) \div (\frac{4}{5})$

For Problems 1 through 10, carry out the indicated operations in the agreed-upon order.

- | | |
|----------------------------------|-----------------------------------|
| 1. $5 + 6 \times 7$ | 2. $3 + 8 \times 7$ |
| 3. $9 - 4 + 5$ | 4. $11 - 6 + 4$ |
| 5. $12 \div 3 \times 2$ | 6. $18 \div 9 \times 2$ |
| 7. $7 - 8 \div 2 + 4$ | 8. $24 - 12 \times 2 + 4$ |
| 9. $16 - 4 + 12 \div 6 \times 2$ | 10. $50 - 30 \times 2 + 8 \div 2$ |

For Problems 11 through 24, evaluate the given expression

- (a) for $x = 2$
 (b) for $x = -3$.

- | | |
|----------------------|---------------------|
| 11. $4x - 1$ | 12. $3x - 5$ |
| 13. $ 3x - 5 $ | 14. $ 4x - 1 $ |
| 15. $5 - 7x - 8$ | 16. $8 - 5x - 2$ |
| 17. $ 8 - 5x - 2$ | 18. $ 5 - 7x - 8$ |
| 19. $x^2 - 4x + 6$ | 20. $x^2 + 6x - 9$ |
| 21. $4x^2 - 5x - 11$ | 22. $5x^2 - 7x + 1$ |
| 23. $5 - 2 \cdot x$ | 24. $3 + 4 \cdot x$ |

For Problems 25 through 40, simplify the given expression.

- | | |
|---|-----------------------------------|
| 25. $6 - [5 - (3 - x)]$ | 26. $2x - [3x + (x - 2)]$ |
| 27. $7(x - 2(3 - x))$ | 28. $3(6x - 5(x - 1))$ |
| 29. $7 - 2[3 - 2(x + 4)]$ | 30. $8 + 4[5 - 6(x - 2)]$ |
| 31. $3x - [2x + (x - 5)]$ | 32. $4x - [3x - (2x - x)]$ |
| 33. $6 - 2[x - 3 - (x + 4) + 3(x - 2)]$ | |
| 34. $7[2 - 3(x - 4) + 4(x - 6)]$ | |
| 35. $6[x - \frac{1}{2}(x - 1)]$ | 36. $8[2x - \frac{1}{4}(6x + 5)]$ |



37. $x^2 + y^2 - [x(x + y) - y(y - x)]$
38. $4x^2 - 2x(x - 2y) + 2y(2y + x) - 2x^2$
39. $-(-(-(-x)))$ 40. $x - [x - (x - \overline{x - y})]$
41. Calvin Butterball and Phoebe Small evaluate the expression $|x - 3|$ for $x = 7$, getting:
- Calvin: $|x - 3| = |7 - 3| = 7 + 3 = \underline{\underline{10}}$
- Phoebe: $|x - 3| = |7 - 3| = |4| = \underline{\underline{4}}$
- Who is right? What mistake did the other one make?
42. Kay Oss evaluates the expression $|x + 2| - 5x$ by substituting 7 for the first x and 3 for the second x . What axiom did Kay violate?

1-4

POLYNOMIALS

Polynomials are algebraic expressions that involve only the operations of *addition*, *subtraction*, and *multiplication* of variables. For example,

$$3x^2 + 5x - 7, \quad x + 2, \quad \text{and} \quad xy^3z^2$$

are polynomials. They involve no non-algebraic operations such as absolute value, and no operations under which the set of real numbers is not closed, such as division and square root. Thus, polynomials stand for *real* numbers no matter what real values you substitute for the variables.

Objectives:

1. Given an expression, tell whether or not it is a polynomial. If it is, then *name* it by “degree” and by number of terms.
2. Given two binomials, multiply them together.

Notes:

1. The expression $\frac{3}{x-3}$ is *not* a polynomial since it involves *division* by a *variable*. If x were 5, the expression would have the form $\frac{3}{2}$, which is *not* a real number.
2. The expression $\sqrt{x}$ is *not* a polynomial since it involves the *square root* of a *variable*. If x were less than 0, the expression would stand for an imaginary number rather than a real number.

3. The expression $|x - 7|$ is *not* a polynomial since it involves the *non-algebraic operation* “absolute value.”
4. Expressions such as $\sqrt{3x}$ and $\frac{x}{3}$ (which equals $\frac{1}{3} \cdot x$) are considered to be polynomials since the operations $\div$ and $\sqrt{\quad}$ are performed on *constants* rather than variables.
5. The operation exponentiation (“raising to powers”) is *not* listed among the polynomial operations. If the exponent is an integer, such as in x^4 , then exponentiation is just repeated multiplication. So expressions with only *integer* exponents are polynomials. In Chapter 6 you will learn what happens when the exponent is not an integer.

“Terms” in an expression are parts of the expression that are *added* or *subtracted*. For example, the expression

$$3x^2 + 5x - 7$$

has three terms, namely, $3x^2$, $5x$, and 7 . Special names are used for expressions that have 1, 2, or 3 terms.

NAMES

No. of Terms	Name	Example
1	monomial	$3x^2y^5$
2	binomial	$3x^2 + y^5$
3	trinomial	$3 - x^2 + y^5$
4 or more	(no special name)	$3x^5 - 2x^4 + 5x^3 - 6x^2 + 2x$

The word “polynomial” originally meant “many terms.” However, it is possible to get a *monomial* by adding two polynomials. For example,

$$(3x^2 + 5x - 7) + (8x^2 - 5x + 7) = 11x^2,$$

a *monomial*. By calling monomials, binomials, and trinomials “polynomials,” too, the set of polynomials has the desirable property of being *closed* under addition. It is also closed under multiplication.

“Factors” in an expression are parts of the expression that are *multiplied* together. For example, $5x^2$ has *three* factors, 5, x , and x . Special names are given to polynomials depending on how many *variables* are *multiplied* together.

For example, $3x^2y^5$ is *seventh* degree because seven variables are multiplied together ($x \cdot x \cdot y \cdot y \cdot y \cdot y \cdot y$). But $3x^2 + y^5$ is only *fifth* degree because at most five variables are multiplied together ($y \cdot y \cdot y \cdot y \cdot y$). An expression such as $17x$ that has only *one* variable is called *first* degree, and a constant such as 17 which has *no* variable is called *zero* degree.



DEFINITION

DEGREE OF A POLYNOMIAL

The **degree** of a polynomial is the maximum number of variables that appear as factors in any one term.

Various degrees are given special names, as follows:

NAMES

Degree	Name	Example	Memory Aid
0	constant	13	Constants do not vary.
1st	linear	$5x$	A line has <i>one</i> dimension.
2nd	quadratic	$7x^2$	A square is a quadrangle.
3rd	cubic	$4x^3$	A cube has <i>three</i> dimensions.
4th	quartic	x^4	A <i>quart</i> is a <i>fourth</i> of a gallon.
5th	quintic	$9x^5$	Quintuplets are <i>five</i> children.
6th or more	(no special name)	$3x^{17}$	(Make up your <i>own</i> names, Hectic, Septic, etc.)

Notes:

- Various parts of a monomial such as $3x^2$ have special names.
 3 is the *numerical coefficient*.
 x is the *base*.
 2 is the *exponent*.
 x^2 is a *power* (the second power of x).
- "Zero" could have *any* degree, because 0 equals $0x^3$, $0x^{15}$, $0x^{1066}$, etc. To avoid this difficulty, 0 is usually called a polynomial with *no* degree.

Multiplying Binomials: Multiplying binomials requires a *double* use of the distributive property.

For example,

$$(x - 3)(2x + 5)$$

can be thought of as

$$\text{number} \times (2x + 5).$$

Distributing the "number," you get

$$\text{number} \times 2x + \text{number} \times 5.$$

Algebra and Trigonometry

Functions and Applications

Special EDITION

- 1 Preliminary Information
- 2 Functions and Relations
- 3 Linear Functions
- 4 Systems of Linear Equations and Inequalities
- 5 Quadratic Functions and Complex Numbers
- 6 Exponential and Logarithmic Functions
- 7 Rational Algebraic Functions
- 8 Irrational Algebraic Functions
- 9 Quadratic Relations and Systems
- 10 Higher-Degree Functions and Complex Numbers
- 11 Sequences and Series
- 12 Probability, Data Analysis, and Functions of a Random Variable
- 13 Trigonometric and Circular Functions
- 14 Properties of Trigonometric and Circular Functions
- 16 Triangle Problems

